arXiv:1301.7352v3 [hep-th] 28 Feb 2013

KUNS-2432
MISC-2013-01

On propagators in de Sitter space

Masafumi Fukuma,[1*] Yuho Sakatani,[2†] and Sotaro Sugishita[1‡]

[1] *Department of Physics, Kyoto University*
Kyoto 606-8502, Japan

[2] *Maskawa Institute for Science and Culture, Kyoto Sangyo University*
Kyoto 603-8555, Japan

Abstract

In a spacetime with no global timelike Killing vector, we do not have a natural choice for the vacuum state of matter fields, leading to the ambiguity in defining the Feynman propagators. In this paper, choosing the vacuum state as the instantaneous ground state of the Hamiltonian at each moment, we develop a method for calculating wave functions associated with the vacuum and the corresponding in-in and in-out propagators. We apply this method to free scalar field theory in de Sitter space and obtain de Sitter invariant propagators in various coordinate patches. We show that the in-out propagator in the Poincaré patch has a finite massless limit with the full de Sitter invariance preserved. We argue and numerically check that our in-out propagators coincide with those obtained by the path integral with the standard iε prescription. We also show that the in-out propagators satisfy Polyakov's composition principle. Several applications of our propagators are also discussed.

---

[*] E-mail address: fukuma_at_gauge.scphys.kyoto-u.ac.jp
[†] E-mail address: yuho_at_cc.kyoto-su.ac.jp
[‡] E-mail address: sotaro_at_gauge.scphys.kyoto-u.ac.jp

Contents

1 Introduction 2

2 General framework 4

2.1 Setup 4
2.2 Quantum mechanics with time-dependent Hamiltonian 6
2.3 Heisenberg picture 8
2.4 Bogoliubov coefficients for finite time intervals 11
2.5 Wave functions 12
2.6 Feynman propagators 14

3 Simple example: scalar field in Minkowski space 16

3.1 Setup 16
3.2 Propagator for each mode 17
3.3 Propagator in spacetime 18

4 Scalar field in de Sitter space 19

4.1 Geometry and definitions 19
4.2 Scalar field in the Poincaré patch 21
  4.2.1 Propagators for each mode in the Poincaré patch 22
  4.2.2 Propagators in the Poincaré patch 25
4.3 Scalar field in the global patch 26
  4.3.1 Propagators for each mode in the global patch 27
  4.3.2 Propagators in the global patch 30

5 Feynman path integral in de Sitter space 33

6 Heat kernel representation and the composition principle 37

6.1 General theory 37
6.2 de Sitter case 40

6.3 Relation to the Green function in Euclidean AdS space 42

7 Discussions and conclusion **44**

A Proof of Eq. (2.69) **46**

B Asymptotically Minkowski space **47**

B.1 Setup 47

B.2 Wave functions 48

C Propagator in Minkowski space **50**

D Proofs of Eqs. (4.47) **and** (4.51) **52**

E Associated Legendre functions and the addition formulae **53**

F Integral representation of the associated Legendre functions **62**

1. Introduction

In a spacetime with no global timelike Killing vector, we do not have an established prescription to define the vacuum state of matter fields, along with which the propagator cannot be defined uniquely. de Sitter space is a typical example of such spacetime, and various vacua have been studied throughout the decades. Among them, the Euclidean vacuum (or the Bunch-Davies vacuum [1]) is de Sitter invariant and satisfies the Hadamard condition. This vacuum is regarded as the most natural vacuum in cosmology and is used to describe the physics in the inflationary era. Other de Sitter invariant vacua, called α-vacua [2, 3], have also been studied frequently ($\alpha = -\infty$ corresponds to the Euclidean vacuum). However, for now, there is no conclusive reason to single out one vacuum.

An interesting feature of de Sitter space is its thermodynamic property. As is pointed out in [4], a particle detector staying in de Sitter space behaves as if it is in a thermal bath with the temperature $T = 1/2\pi\ell$, where ℓ is the de Sitter radius. The calculation crucially depends on the choice of the vacuum at the initial time when the detector starts its operation. For example, the α-vacua do not predict the thermal behavior for $\alpha \neq -\infty$ [5]. In this sense, the choice of the vacuum is also important in understanding the thermodynamic behavior of curved spacetimes.

In this paper, we define the vacuum of a free scalar field as the instantaneous ground state of the Hamiltonian at each moment. We develop a method to calculate transition amplitudes during finite time intervals for quantum mechanics with time-dependent Hamiltonian, and define the propagators as the limit of two-point functions when the initial and final times are sent to the past and future infinities.[1] In our method, wave functions associated with the vacuum are automatically determined with no need to consider asymptotic boundary conditions such as positive-energy conditions.

We apply the method to construct various Feynman propagators in de Sitter space. We treat the principal ($m > (d-1)/2$) and the complementary series ($m < (d-1)/2$) at the same time, and show that the obtained in-in and in-out propagators always take de Sitter invariant forms. Furthermore, we show that our de Sitter invariant in-out propagator has a finite massless limit in the Poincaré patch.[2] This is in contrast to the in-in propagators, for which the no-go theorem is known that there is no de Sitter invariant Fock vacuum for massless scalar fields [3].

We argue and numerically check that our in-out propagators coincide with the propagators obtained by the path integral with the standard $\mathrm{i}\varepsilon$ prescription. This result is consistent with the de Sitter invariance of the propagators since the corresponding path integral is performed over a patch which is invariant under the action of de Sitter group. Moreover, our in-out propagators are shown to satisfy the composition principle [9], which has been claimed by Polyakov recently as the property to be satisfied in order for the propagator to be interpreted as representing the sum over paths of a particle moving in a spacetime.

This paper is organized as follows. In section 2, we develop a general framework to calculate the wave functions and propagators for quantum mechanics with time-dependent Hamiltonian. In section 3 we demonstrate how our prescription works in the simplest spacetime, Minkowski space. Another well-studied example of asymptotically Minkowski space is investigated in Appendix B. We then analyze the de Sitter case in section 4. After giving a brief review on the geometry of de Sitter space in subsection 4.1, we discuss the propagators in the Poincaré patch in subsection 4.2 and those in the global patch in subsection 4.3. For the both patches, we first make a mode expansion of a scalar field and calculate the propagators for each mode. We then make the summation over modes to obtain the propagators in spacetime, both of the in-in and in-out types. The obtained propagators are found to be invariant under the full de Sitter group. In section 5 we argue and give an evidence based on numerical calculation that our in-out propagators coincide with those obtained by the path integral with the standard iε prescription. In section 6 we prove that our in-out propagators have the heat-kernel representation, which means that the propagators satisfy Polyakov's composition principle [10]. Section 7 is devoted to discussions and conclusion. We collect miscellaneous formulae in appendices.

---

[1] There had been a study to define the vacuum as the instantaneous ground state, which is sometimes called the *instantaneous Hamiltonian diagonalization method* (see, e.g., [6, 7, 8] and references therein). Our framework has the same principle as that of the method in determining the vacuum, but has an advantage over the method in that it can determine an explicit form of various propagators for finite time intervals and can be applied for a wide class of nonstatic spacetimes.

[2] The in-out propagator in the global patch still diverges in the massless limit.

2. General framework

2.1. Setup

In this paper, we consider quantum theory of a free scalar field $\phi(x)$ living in a d-dimensional curved spacetime with background metric $g_{\mu\nu}$:[3]

$$S[\phi(x)] = -\frac{1}{2}\int \mathrm{d}^d x\,\sqrt{-g}\,\big(g^{\mu\nu}\,\partial_\mu\phi\,\partial_\nu\phi + m^2\,\phi^2\big)\,. \tag{2.1}$$

We assume that the spacetime is globally hyperbolic and the foliation of spacetime (i.e., the set of timeslices) is already specified. We denote the temporal and spatial coordinates by t and $\boldsymbol{x}$, respectively, and the spacetime coordinates by $x = (x^\mu) = (t, \boldsymbol{x})$ ($\mu = 0, 1, \ldots, d-1$). We further assume that the metric has the following form:

$$\mathrm{d}s^2 = -N^2(t)\,\mathrm{d}t^2 + A^2(t)\,h_{ij}(\boldsymbol{x})\,\mathrm{d}x^i\,\mathrm{d}x^j \quad (i, j = 1, \ldots, d-1)\,. \tag{2.2}$$

The action is then written as

$$S = \frac{1}{2}\int \mathrm{d}t \int \mathrm{d}^{d-1}x\,\sqrt{h}\,N\,A^{d-1}\Big(N^{-2}\,\partial_t\phi\,\partial_t\phi + A^{-2}\,\phi\,\Delta_{d-1}\phi - m^2\phi^2\Big)\,, \tag{2.3}$$

where $\sqrt{h} \equiv \sqrt{\det h_{ij}}$, and $\Delta_{d-1} \equiv (1/\sqrt{h})\,\partial_i\big(\sqrt{h}\,h^{ij}\,\partial_j\big)$ is the Laplacian for the spatial metric $\mathrm{d}s^2_{d-1} = h_{ij}(\boldsymbol{x})\,\mathrm{d}x^i\,\mathrm{d}x^j$. We have neglected the surface term coming from integration by parts. We introduce a complete system $\{Y_n(\boldsymbol{x})\}$ of orthonormal eigenfunctions of Δ_{d-1} satisfying

---

[3] The metric has the signature $(-, +, \ldots, +)$.

$$\Delta_{d-1}\,Y_n(\boldsymbol{x}) = -\,\lambda_n\,Y_n(\boldsymbol{x})\,, \qquad \int \mathrm{d}^{d-1}\boldsymbol{x}\,\sqrt{h(\boldsymbol{x})}\,Y_n(\boldsymbol{x})\,Y_{n'}(\boldsymbol{x}) = \delta_{nn'}\,, \tag{2.4}$$

and make the mode expansion of the scalar field as

$$\phi(x) = \phi(t,\boldsymbol{x}) = \sum_n \phi_n(t)\,Y_n(\boldsymbol{x})\,. \tag{2.5}$$

The action can then be written as the sum of the actions for mode functions $\{\phi_n(t)\}$:

$$S = \sum_n \int \mathrm{d}t\,L_n\big(\phi_n(t),\dot{\phi}_n(t),t\big) \tag{2.6}$$

with

$$L_n(\phi_n,\dot{\phi}_n,t) = \frac{\rho(t)}{2}\,\dot{\phi}_n^2 - \frac{\rho(t)\,\omega_n^2(t)}{2}\,\phi_n^2\,, \tag{2.7}$$

$$\rho(t) \equiv N^{-1}(t)\,A^{d-1}(t)\,, \qquad \omega_n(t) \equiv N(t)\,\sqrt{\lambda_n A^{-2}(t) + m^2}\,. \tag{2.8}$$

This shows that the n^{th} mode function $\phi_n(t)$ behaves as a quantum oscillator with time-dependent mass $\rho(t)$ and frequency $\omega_n(t)$, and is described by the following time-dependent Hamiltonian in the Schrödinger picture:

$$H_{n,\,s}(t) = H_n\big(\phi_{n,\,s},\,\pi_{n,\,s},\,t\big) = \frac{1}{2\,\rho(t)}\,\pi_{n,\,s}^2 + \frac{\rho(t)\,\omega^2(t)}{2}\,\phi_{n,\,s}^2\,, \tag{2.9}$$

where the suffix s indicates that the operators are in the Schrödinger picture. Thus, the theory is reduced to quantum mechanics of a set of independent harmonic oscillators with time-dependent parameters:

$$H_s(t) = \sum_n H_{n,\,s}(t) = \sum_n \Big[\frac{1}{2\,\rho(t)}\,\pi_{n,\,s}^2 + \frac{\rho(t)\,\omega^2(t)}{2}\,\phi_{n,\,s}^2\Big]\,. \tag{2.10}$$

The quantization is accomplished by setting the commutation relations

$$[\phi_{n,\,s},\,\pi_{m,\,s}] = \mathrm{i}\,\delta_{n,m}\,, \quad [\phi_{n,\,s},\,\phi_{m,\,s}] = 0 = [\pi_{n,\,s},\,\pi_{m,\,s}]\,. \tag{2.11}$$

In the following subsections, we develop a general theory to describe the time evolution of states for quantum mechanics with such a time-dependent Hamiltonian.

We here make a comment on a subtlety existing in field redefinitions. By transforming the mode function $\phi_n(t)$ to $\chi_n(t) = \rho^{-1/2}(t)\,\phi_n(t) \equiv e^{\sigma(t)}\,\phi_n(t)$, one can make the coefficient of the kinetic term to unity:

$$S[\chi_n(t)] = \int_{t_i}^{t_f}\mathrm{d}t\,\frac{1}{2}\,\big[\dot{\chi}_n^2(t) - \Omega_n^2(t)\,\chi_n^2(t)\big]\,, \tag{2.12}$$

$$\Omega_n^2(t) \equiv \omega_n^2(t) - \big(\dot{\sigma}(t)\big)^2 - \ddot{\sigma}(t)\,. \tag{2.13}$$

However, it can often happen that $\Omega_n^2(t)$ takes a negative value for some region of m^2 even though the original $\omega^2(t)$ is strictly positive.[4] Although the physics should be the same for the two descriptions using $\phi_n(t)$ and $\chi_n(t)$, the inverted harmonic potential for $\chi_n(t)$ can easily cause a catastrophe when making an analysis based on an approximation such as the WKB approximation. In order to avoid this subtlety (and also to keep the original symmetry manifest), we will not make such transformations.[5]

2.2. Quantum mechanics with time-dependent Hamiltonian

For brevity, in the following discussions, we omit the mode index n and denote the canonical variables $\{\phi_{n,s},\ \pi_{n,s}\}$ in the Schrödinger picture simply by $\{q_s, p_s\}$. Our Hamiltonian then takes the form

$$H_s(t) = H(q_s, p_s, t) = \frac{1}{2\rho(t)}\, p_s^2 + \frac{\rho(t)\,\omega^2(t)}{2}\, q_s^2\,, \tag{2.14}$$

and the system is quantized by setting the commutation relation

$$[q_s, p_s] = \mathrm{i}\,. \tag{2.15}$$

As we will see below, in order to discuss the behavior of states near the temporal boundary in a well-defined manner, we need to let the frequency $\omega(t)$ have a small negative imaginary part without breaking the symmetry existing in the background spacetime. This can be best realized by replacing m^2 in (2.1) by $m^2 - \mathrm{i}\varepsilon$ ($\varepsilon > 0$). Note that the Hamiltonian $H_s(t)$ is no longer hermitian when $\varepsilon \neq 0$.

We denote by T_s the time at which quantization is carried out in the Schrödinger picture. The Hilbert state $\mathcal{H} = \{|\psi\rangle\}$ with the hermitian inner product (ψ_1, ψ_2) is then constructed on the timeslice at $t = T_s$, and the dual space $\mathcal{H}^* = \{\langle\psi|\}$ is defined with respect to the hermitian inner product with the rule $\langle\psi_1| \equiv \big(|\psi_1\rangle\big)^\dagger$, i.e., $\langle\psi_1|(|\psi_2\rangle) = (\psi_1, \psi_2)$. The time evolution of a state $|\psi\rangle \in \mathcal{H}$ is governed by the Schrödinger equation

$$\partial_t\,|\psi, t\rangle = -\mathrm{i} H_s(t)\,|\psi, t\rangle \tag{2.16}$$

---

[4] A typical example is a scalar field in the Poincaré patch of de Sitter space. One can easily see that $\Omega_n^2(t)$ can be negative when the mass is small and $\phi_n(t)$ represents a mode of long wave length.

[5] The coefficient of the kinetic term can be set to unity also by making a transformation of the time coordinate. We will see that physical quantities do not change under the transformation (see the last paragraph of subsec. 2.4).

with the initial condition $|\psi, T_s\rangle = |\psi\rangle$. The Schrödinger equation can be integrated to the form

$$|\psi, t\rangle = U(t, T_s)\,|\psi\rangle\,, \tag{2.17}$$

where $U(t, T_s)$ is the time-evolution operator expressed as the time-ordered exponential of $H_s(t)$,

$$\begin{aligned} U(t, T_s) &\equiv \mathrm{T}\exp\Bigl(-\mathrm{i}\int_{T_s}^{t} \mathrm{d}t'\, H_s(t')\Bigr) \\ &\equiv \lim_{\Delta t_k \to 0} \bigl(1 - \mathrm{i}\Delta t_N H_s(t_N)\bigr)\bigl(1 - \mathrm{i}\Delta t_{N-1} H_s(t_{N-1})\bigr)\cdots\bigl(1 - \mathrm{i}\Delta t_1 H_s(t_1)\bigr) \\ &\qquad \bigl(t = t_N > t_{N-1} > \cdots > t_1 > t_0 = T_s\,;\ \Delta t_k = t_k - t_{k-1}\bigr)\,. \end{aligned} \tag{2.18}$$

The hermitian conjugate of $|\psi, t\rangle$ is given by

$$\langle\psi, t| = \langle\psi|\, U^\dagger(t, T_s)\,. \tag{2.19}$$

In addition, we introduce a one-parameter family of states for a given state $\overline{\langle\psi|} \in \mathcal{H}^*$ as

$$\overline{\langle\psi, t|} \equiv \overline{\langle\psi|}\, U^{-1}(t, T_s)\,, \tag{2.20}$$

which satisfy

$$\partial_t\, \overline{\langle\psi, t|} = +\mathrm{i}\, \overline{\langle\psi, t|}\, H_s(t)\,, \quad \overline{\langle\psi, T_s|} = \overline{\langle\psi|}\,. \tag{2.21}$$

Note that the paring of $\overline{\langle\psi_1|}$ and $|\psi_2\rangle$ does not change under the time evolution,

$$\overline{\langle\psi_1, t|}\psi_2, t\rangle = \overline{\langle\psi_1|}\psi_2\rangle\,, \tag{2.22}$$

although this is not the case for $\langle\psi_1, t|\psi_2, t\rangle$ when $\omega(t)$ has a nonvanishing imaginary part because the time evolution operator is then not unitary, $U^{-1}(t, T_s) \neq U^\dagger(t, T_s)$.

The spectrum of the Hamiltonian $H_s(t)$ can be easily found by introducing, as usual, a pair of operators,[6]

$$a_s(t) \equiv \sqrt{\frac{\rho(t)\,\omega(t)}{2}}\, q_s + \mathrm{i}\sqrt{\frac{1}{2\rho(t)\,\omega(t)}}\, p_s\,, \tag{2.23}$$

$$\bar{a}_s(t) \equiv \sqrt{\frac{\rho(t)\,\omega(t)}{2}}\, q_s - \mathrm{i}\sqrt{\frac{1}{2\rho(t)\,\omega(t)}}\, p_s\,. \tag{2.24}$$

---

[6] They are Schrödinger operators and the time dependence comes only through the parameters $\rho(t)$ and $\omega(t)$.

We call $a_s(t)$ and $\bar{a}_s(t)$ the annihilation and creation operators at time t. Note that $\bar{a}_s(t) \neq a_s^\dagger(t)$ when $\omega(t)$ has a nonvanishing imaginary part. From the commutation relation (2.15), we have

$$[a_s(t), \bar{a}_s(t)] = 1\,. \tag{2.25}$$

The Hamiltonian (2.14) can then be rewritten as

$$H_s(t) = \frac{\omega(t)}{2}\left[\bar{a}_s(t)\,a_s(t) + a_s(t)\,\bar{a}_s(t)\right] = \omega(t)\left[\bar{a}_s(t)\,a_s(t) + \frac{1}{2}\right]. \tag{2.26}$$

We define the state $|0_t, t\rangle$ as that which vanishes when acted on by $a_s(t)$:

$$a_s(t)\,|0_t, t\rangle = 0\,. \tag{2.27}$$

Similarly, the state $\overline{\langle 0_t, t|}$ is defined as[7]

$$\overline{\langle 0_t, t|}\,\bar{a}_s(t) = 0\,. \tag{2.28}$$

Then the left and right eigenstates of $H_s(t)$ are given by

$${}_t\overline{\langle n, t|} \equiv \frac{1}{n!}\,\overline{\langle 0_t, t|}\left[a_s(t)\right]^n, \quad |n, t\rangle_t \equiv \frac{1}{n!}\left[\bar{a}_s(t)\right]^n |0_t, t\rangle, \tag{2.29}$$

which satisfy

$${}_t\overline{\langle n, t|}\,H_s(t) = \left(n + \frac{1}{2}\right)\omega(t)\,{}_t\overline{\langle n, t|}\,, \quad H_s(t)\,|n, t\rangle_t = \left(n + \frac{1}{2}\right)\omega(t)\,|n, t\rangle_t\,. \tag{2.30}$$

We call $|0_t, t\rangle = |0, t\rangle_t$ the ground state (or the vacuum) at time t, since this is the minimum energy state when $\varepsilon = 0$.

It is important to note that the ground state at time t', $|0_{t'}, t'\rangle$, is generically different from the state $|0_t, t'\rangle$; the latter is obtained as the time evolution of the ground state $|0_t, t\rangle$ at time t, $|0_t, t'\rangle = U(t', t)|0_t, t\rangle$ (see Fig. 1). Since the Hamiltonian is already specified at each time, the vacuum state is uniquely determined, and there is no freedom to introduce other vacuum states through Bogoliubov transformations without changing the timeslices (or Hamiltonian).

2.3. Heisenberg picture

Now we move from the Schrödinger picture to the Heisenberg picture. Given a Schrödinger operator $O_s(t)$ [possibly depending on t through the parameters involved when constructing

---

[7] Note that $\overline{\langle 0_t|} = \overline{\langle 0_t, T_s|}$ may differ from $\langle 0_t| = \langle 0_t, T_s|$ since $\bar{a}_s(T_s) \neq a^\dagger(T_s)$ when $\mathrm{Im}\,\omega(t) \neq 0$.



Figure 1: Time evolution of states. The system is quantized in the Schrödinger picture on the timeslice at T_s . $|0_t, t\rangle$ is the state annihilated by the Schrödinger operator $a_s(t)$, while $|0_t\rangle$ is the state annihilated by the Heisenberg operator $a(t)$.

the operator as in Eq. (2.23)], we define the corresponding Heisenberg operator as

$$O(t) \equiv U^{-1}(t, T_s)\, O_s(t)\, U(t, T_s)\,, \tag{2.31}$$

which satisfies the Heisenberg equation,

$$\dot{O}(t) = \mathrm{i}\,[H(t),\, O(t)] + \frac{\partial O(t)}{\partial t} \quad \Big(\frac{\partial O(t)}{\partial t} \equiv U^{-1}(t, T_s)\,\frac{\partial O_s(t)}{\partial t}\,U(t, T_s)\Big)\,. \tag{2.32}$$

For our harmonic oscillator, the time evolution of the canonical variables is given by

$$\dot{q}(t) = \mathrm{i}\,[H(t),\, q(t)] = \frac{p(t)}{\rho(t)}\,, \tag{2.33}$$

$$\dot{p}(t) = \mathrm{i}\,[H(t),\, p(t)] = \,-\,\rho(t)\,\omega^2(t)\,q(t)\,, \tag{2.34}$$

and by eliminating $p(t)$, we obtain the differential equation

$$\frac{\mathrm{d}}{\mathrm{d}t}\Big(\rho(t)\,\frac{\mathrm{d}}{\mathrm{d}t}\,q(t)\Big) + \rho(t)\,\omega^2(t)\,q(t) = 0\,. \tag{2.35}$$

This is certainly the equation of motion derived from the Lagrangian $L(q, \dot{q}, t) = \rho(t)\,\dot{q}^2/2 - \rho(t)\,\omega^2(t)\,q^2/2$. Note that if we had used $U^\dagger(t, T_s)$ in Eq. (2.31) instead of $U^{-1}(t, T_s)$, the equation of motion could not be reproduced correctly when $\omega(t)$ has an imaginary part.

In the Heisenberg picture, the annihilation and creation operators become

$$a(t) \equiv U^{-1}(t,T_s)\,a_s(t)\,U(t,T_s) = \sqrt{\frac{\rho(t)\,\omega(t)}{2}}\,q(t) + \mathrm{i}\sqrt{\frac{1}{2\rho(t)\,\omega(t)}}\,p(t)\,, \tag{2.36}$$

$$\bar{a}(t) \equiv U^{-1}(t,T_s)\,\bar{a}_s(t)\,U(t,T_s) = \sqrt{\frac{\rho(t)\,\omega(t)}{2}}\,q(t) - \mathrm{i}\sqrt{\frac{1}{2\rho(t)\,\omega(t)}}\,p(t)\,, \tag{2.37}$$

and satisfy the commutation relation

$$[a(t),\bar{a}(t)] = 1\,. \tag{2.38}$$

Note that they satisfy the following equations:

$$a(t)\,|0_t\rangle = 0 = \langle 0_t|\,a^\dagger(t)\,, \qquad \overline{\langle 0_t|}\,\bar{a}(t) = 0 = \bar{a}^\dagger(t)\,\overline{|0_t\rangle}\,. \tag{2.39}$$

The Hamiltonian is then expressed as

$$H(t) \equiv U^{-1}(t,T_s)\,H_s(t)\,U(t,T_s) = \frac{p^2(t)}{2\rho(t)} + \frac{\rho(t)\,\omega^2(t)\,q^2(t)}{2} = \omega(t)\left[\bar{a}(t)\,a(t) + \frac{1}{2}\right]. \tag{2.40}$$

We denote by $\{f(t),g(t)\}$ a pair of linearly independent c-number solutions of (2.35). One can easily show that their Wronskian,

$$W[f,g](t) \equiv f(t)\,\dot{g}(t) - \dot{f}(t)\,g(t)\,, \tag{2.41}$$

satisfies the equation

$$\frac{\mathrm{d}}{\mathrm{d}t}(\rho\,W[f,g]) = f\,\frac{\mathrm{d}}{\mathrm{d}t}(\rho\,\dot{g}) - \frac{\mathrm{d}}{\mathrm{d}t}(\rho\,\dot{f})\,g = 0\,. \tag{2.42}$$

Thus the combination (to be called the weighted Wronskian)

$$W_\rho[f,g] \equiv \rho(t)\,W[f,g](t) \tag{2.43}$$

does not depend on t. Since $q(t)$ is also a solution of Eq. (2.35), we can expand canonical variables $q(t)$ and $p(t)$ as

$$q(t) = c_1\,f(t) + c_2\,g(t)\,, \tag{2.44}$$

$$p(t) = \rho(t)\,\dot{q}(t) = \rho(t)\,\big[\,c_1\,\dot{f}(t) + c_2\,\dot{g}(t)\,\big]\,, \tag{2.45}$$

where c_1 and c_2 are some time-independent quantum operators. Equations (2.44) and (2.45) can be solved with respect to c_1 and c_2 as

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \frac{1}{W_\rho[f,g]}\begin{pmatrix} \rho\,\dot{g} & -g \\ -\rho\,\dot{f} & f \end{pmatrix}(t)\begin{pmatrix} q(t) \\ p(t) \end{pmatrix}\,. \tag{2.46}$$

This can be further rewritten by using (2.36) and (2.37) as

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = C(t) \begin{pmatrix} a(t) \\ \bar{a}(t) \end{pmatrix} , \tag{2.47}$$

where

$$C(t) \equiv \frac{1}{W_\rho[f,g]\,\sqrt{2\rho(t)\,\omega(t)}} \begin{pmatrix} v(t) & \bar{v}(t) \\ -u(t) & -\bar{u}(t) \end{pmatrix} \tag{2.48}$$

with

$$\left\{ \begin{array}{c} u(t) \\ \bar{u}(t) \end{array} \right\} \equiv \rho(t)\,\big[\,\dot{f}(t) \pm \mathrm{i}\omega(t)\,f(t)\,\big] , \quad \left\{ \begin{array}{c} v(t) \\ \bar{v}(t) \end{array} \right\} \equiv \rho(t)\,\big[\,\dot{g}(t) \pm \mathrm{i}\omega(t)\,g(t)\,\big] . \tag{2.49}$$

Note that

$$\big(u\,\bar{v} - v\,\bar{u}\big)(t) = 2\mathrm{i}\rho(t)\,\omega(t)\,W_\rho[f,g] , \tag{2.50}$$

$$\det C(t) = \frac{\mathrm{i}}{W_\rho[f,g]} \quad \big(= \mathrm{const}\big) , \tag{2.51}$$

$$C^{-1}(t) = \frac{-\mathrm{i}}{\sqrt{2\rho(t)\,\omega(t)}} \begin{pmatrix} -\bar{u}(t) & -\bar{v}(t) \\ u(t) & v(t) \end{pmatrix} . \tag{2.52}$$

2.4. Bogoliubov coefficients for finite time intervals

The Bogoliubov coefficients from time t' to time t are defined by

$$\begin{pmatrix} a(t) \\ \bar{a}(t) \end{pmatrix} \equiv \begin{pmatrix} \bar{\alpha} & -\bar{\beta} \\ -\beta & \alpha \end{pmatrix} (t;t') \begin{pmatrix} a(t') \\ \bar{a}(t') \end{pmatrix} . \tag{2.53}$$

Since the operators $\{c_1, c_2\}$ in (2.47) do not depend on time, we have

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = C(t) \begin{pmatrix} a(t) \\ \bar{a}(t) \end{pmatrix} = C(t') \begin{pmatrix} a(t') \\ \bar{a}(t') \end{pmatrix} , \tag{2.54}$$

from which we find

$$\begin{aligned} &\begin{pmatrix} \bar{\alpha} & -\bar{\beta} \\ -\beta & \alpha \end{pmatrix} (t;t') = C^{-1}(t)\,C(t') \\ &= \frac{-\mathrm{i}}{2W_\rho[f,g]\,\sqrt{\rho(t)\,\omega(t)\,\rho(t')\,\omega(t')}} \begin{pmatrix} u(t')\,\bar{v}(t) - v(t')\,\bar{u}(t) & \bar{u}(t')\,\bar{v}(t) - \bar{v}(t')\,\bar{u}(t) \\ v(t')\,u(t) - u(t')\,v(t) & \bar{v}(t')\,u(t) - \bar{u}(t')\,v(t) \end{pmatrix} . \end{aligned} \tag{2.55}$$

This is the fundamental formula to express the Bogoliubov coefficients in terms of a given set of independent solutions $\{f(t),\, g(t)\}$.

Due to the commutation relations (2.38), the Bogoliubov coefficients should satisfy the relation

$$\big(\bar{\alpha}\,\alpha - \bar{\beta}\,\beta\big)(t;t') = 1\,. \tag{2.56}$$

This can be directly checked by using the identity (2.51) as

$$(\bar{\alpha}\alpha - \bar{\beta}\beta)(t;t') = \det C^{-1}(t)\,\det C(t') = \frac{W_\rho[f,g]}{W_\rho[f,g]} = 1\,. \tag{2.57}$$

Note that $\bar{\alpha} \neq \alpha^*$ and $\bar{\beta} \neq \beta^*$ when $\omega(t)$ has a nonvanishing imaginary part.

2.5. Wave functions

Using the Bogoliubov coefficients, we can express the Heisenberg operators $q(t)$ with the creation and annihilation operators at a different time t_I as follows:[8]

$$\begin{aligned} q(t) &= \frac{1}{\sqrt{2\rho(t)\,\omega(t)}}\,\big(a(t) + \bar{a}(t)\big) \\ &= \frac{1}{\sqrt{2\rho(t)\,\omega(t)}}\big(\bar{\alpha}(t;t_I)\,a_I - \bar{\beta}(t;t_I)\,\bar{a}_I + \alpha(t;t_I)\,\bar{a}_I - \beta(t;t_I)\,a_I\big) \\ &\equiv \varphi(t;t_I)\,a_I + \bar{\varphi}(t;t_I)\,\bar{a}_I\,, \end{aligned} \tag{2.58}$$

where we have defined functions $\varphi(t;t_I)$ and $\bar{\varphi}(t;t_I)$ (to be called *wave functions*) as

$$\varphi(t;t_I) \equiv \frac{1}{\sqrt{2\rho(t)\,\omega(t)}}\,\big(\bar{\alpha}(t;t_I) - \beta(t;t_I)\big) = \frac{1}{W_\rho[f,g]\,\sqrt{2\rho_I\,\omega_I}}\big(v_I\,f(t) - u_I\,g(t)\big)\,, \tag{2.59}$$

$$\bar{\varphi}(t;t_I) \equiv \frac{1}{\sqrt{2\rho(t)\,\omega(t)}}\,\big(\alpha(t;t_I) - \bar{\beta}(t;t_I)\big) = \frac{1}{W_\rho[f,g]\,\sqrt{2\rho_I\,\omega_I}}\big(\bar{v}_I\,f(t) - \bar{u}_I\,g(t)\big)\,. \tag{2.60}$$

By using the t-independence of $W_\rho[f,g]$ and Eq. (2.50), we can show that $\varphi(t;t_I)$ and $\bar{\varphi}(t;t_I)$ are normalized as

$$W_\rho[\varphi(t;t_I), \bar{\varphi}(t;t_I)] = \mathrm{i} \quad \big(\forall t\,,\ \forall t_I\big)\,. \tag{2.61}$$

Moreover, the following relation holds for any value of t:

$$\begin{pmatrix} \bar{\alpha} & -\bar{\beta} \\ -\beta & \alpha \end{pmatrix}(t_1;t_0) = -\mathrm{i}\begin{pmatrix} -W_\rho[\bar{\varphi}(t;t_1), \varphi(t;t_0)] & -W_\rho[\bar{\varphi}(t;t_1), \bar{\varphi}(t;t_0)] \\ W_\rho[\varphi(t;t_1), \varphi(t;t_0)] & W_\rho[\varphi(t;t_1), \bar{\varphi}(t;t_0)] \end{pmatrix}\,. \tag{2.62}$$

---

[8] In the following, we will use the shorthand notation such as $f_I \equiv f(t_I)$ or $\dot{f}_I \equiv \dot{f}(t_I)$ when a quantity is evaluated at time t_I.

We are now in a position to make a few comments.

basis independence

We can show that the Bogoliubov coefficients and the wave functions $\{\varphi(t;t_I),\bar{\varphi}(t;t_I)\}$ do no depend on the choice of a pair of independent solutions $\{f(t),g(t)\}$, as they should. In fact, suppose that we take another pair $\{f'(t),g'(t)\}$. They should be expressed as linear combinations of $\{f(t),g(t)\}$ of the form

$$\big(f'(t)\ \ g'(t)\big) = \big(f(t)\ \ g(t)\big)\,\Xi \qquad (\Xi \in GL(2,\mathbb{C}))\,, \tag{2.63}$$

from which we have

$$\begin{pmatrix} f' & g' \\ \dot{f}' & \dot{g}' \end{pmatrix}(t) = \begin{pmatrix} f & g \\ \dot{f} & \dot{g} \end{pmatrix}(t)\ \Xi\,, \qquad C'(t) = \Xi^{-1}\,C(t)\,. \tag{2.64}$$

The new Bogoliubov coefficients associated with the choice $\{f'(t),g'(t)\}$ then become

$$\begin{aligned} \begin{pmatrix} \bar{\alpha}' & -\bar{\beta}' \\ -\beta' & \alpha' \end{pmatrix}(t;t_I) &= \big[C'(t)\big]^{-1}\,C'(t_I) = C^{-1}(t)\,\Xi\,\Xi^{-1}\,C(t_I) \\ &= C^{-1}(t)\,C(t_I) = \begin{pmatrix} \bar{\alpha} & -\bar{\beta} \\ -\beta & \alpha \end{pmatrix}(t;t_I)\,, \end{aligned} \tag{2.65}$$

which shows the basis-independence of the Bogoliubov coefficients. The wave functions $\{\varphi(t;t_I),\bar{\varphi}(t;t_I)\}$ are also basis independent since they are expressed by the basis-independent Bogoliubov coefficients [see Eqs. (2.59) and (2.60)].

lapse independence

We can show that the Bogoliubov coefficients and the wave functions $\varphi(t;t_I)$ behave as scalar functions under the temporal reparametrizations preserving the foliation of spacetime. In fact, for such reparametrization $t \to \tilde{t} = \tilde{t}(t)$, the pull-back of the lapse function $N(t)$ [see Eq. (2.2)] is given by $N(t) \to \tilde{N}(t) = (\mathrm{d}\tilde{t}/\mathrm{d}t)\,N\big(\tilde{t}(t)\big)$, and we can choose a new pair of solutions $\{\tilde{f}(t),\tilde{g}(t)\}$ as $\tilde{f}(t) = f\big(\tilde{t}(t)\big)$ and $\tilde{g}(t) = g\big(\tilde{t}(t)\big)$. Then, we can easily show that the functions $\rho(t)\,\omega(t)$, $u(t)$, $v(t)$, and $W_\rho[f(t),g(t)]$ transform as scalar functions under the reparametrization. Since the Bogoliubov coefficients and the wave function $\varphi(t;t_I)$ are written as combinations of these functions, they also transform as scalar functions. This means that there is no need to care about the temporal reparametrization [i.e. the choice of the lapse function $N(t)$] when we construct vacua.

2.6. Feynman propagators

We consider the region $t_i < t_0 \leq \{t, t'\} \leq t_1 < t_f$, where t_f and t_i are the future and the past boundaries of the spacetime region we consider. The in-in and in-out propagators are defined with the following two steps.

step 1

We first introduce the following two-point functions from our wave functions $\varphi(t; t_I)$ and $\bar{\varphi}(t; t_I)$:[9]

$$\begin{aligned} G_{00}(t, t'; t_0, t_0) &\equiv \frac{\langle 0_{t_0} |\, q^\dagger(t_>)\, q(t_<)\, | 0_{t_0} \rangle}{\langle 0_{t_0} | 0_{t_0} \rangle} \\ &= \frac{\mathrm{i}}{W_\rho[\bar{\varphi}^*(t; t_0),\ \bar{\varphi}(t; t_0)](T_s)}\, \bar{\varphi}^*(t_>; t_0)\, \bar{\varphi}(t_<; t_0)\,, \end{aligned} \tag{2.66}$$

$$\begin{aligned} G_{10}(t, t'; t_1, t_0) &\equiv \frac{\overline{\langle 0_{t_1} |}\, \mathrm{T}\, q(t)\, q(t')\, | 0_{t_0} \rangle}{\overline{\langle 0_{t_1} |} 0_{t_0} \rangle} \\ &= \frac{\mathrm{i}}{W_\rho[\varphi(t; t_1), \bar{\varphi}(t; t_0)]}\, \varphi(t_>; t_1)\, \bar{\varphi}(t_<; t_0) \quad (t\text{: arbitrary})\,, \end{aligned} \tag{2.67}$$

where $t_> \equiv \max(t, t')$ and $t_< \equiv \min(t, t')$.

step 2

We then define the in-in and in-out propagators by sending t_0 and t_1 to the values at the temporal boundary:

$$G^{\mathrm{in/in}}(t, t') \equiv \lim_{t_0 \to t_i} G_{00}(t, t'; t_0, t_0)\,, \qquad G^{\mathrm{in/out}}(t, t') \equiv \lim_{\substack{t_0 \to t_i \\ t_1 \to t_f}} G_{10}(t, t'; t_1, t_0)\,. \tag{2.68}$$

The last expression of (2.66) is obtained from the identities

$$\bar{a}_0^\dagger = -\frac{W_\rho[\varphi^*(t; t_0), \bar{\varphi}(t; t_0)](T_s)}{W_\rho[\bar{\varphi}^*(t; t_0), \bar{\varphi}(t; t_0)](T_s)}\, a_0^\dagger + \frac{\mathrm{i}}{W_\rho[\bar{\varphi}^*(t; t_0), \bar{\varphi}(t; t_0)](T_s)}\, a_0\,, \tag{2.69}$$

$$\frac{\langle 0_{t_0} |\, \bar{a}_0^\dagger\, \bar{a}_0\, | 0_{t_0} \rangle}{\langle 0_{t_0} | 0_{t_0} \rangle} = \frac{\mathrm{i}}{W_\rho[\bar{\varphi}^*(t; t_0)\, \bar{\varphi}(t; t_0)](T_s)}\,, \tag{2.70}$$

---

[9] If we instead use $G'_{10}(t, t'; t_1, t_0) = \langle 0_{t_1} |\, q^\dagger(t_>)\, q(t_<)\, | 0_{t_0} \rangle / \langle 0_{t_1} | 0_{t_0} \rangle$, then the corresponding in-out propagator will not coincide with the propagator obtained by the standard path integral (see section 5), and thus we do not consider this choice in this paper. By contrast, the in-in propagator still has options for its definition (e.g., $G'_{00}(t, t'; t_0, t_0) = \langle 0_{t_0} |\, \mathrm{T}\, q(t)\, q(t')\, | 0_{t_0} \rangle / \langle 0_{t_0} | 0_{t_0} \rangle$), and we leave as a future work the detailed study of such options as well as the relation to the path integral based on the Schwinger-Keldysh formalism [11, 12].

which can be shown by using the hermiticity at the time T_s, $q^\dagger(T_s) = q(T_s)$ and $p^\dagger(T_s) = p(T_s)$ (see Appendix A). We then have

$$
\begin{aligned}
G_{00}(t,t';t_0,t_0) &= \frac{\langle 0_{t_0}|\,\big(\varphi(t_>;t_0)\,a_0 + \bar\varphi(t_>;t_0)\,\bar a_0\big)^\dagger\,\big(\varphi(t_<;t_0)\,a_0 + \bar\varphi(t_<;t_0)\,\bar a_0\big)\,|0_{t_0}\rangle}{\langle 0_{t_0}|0_{t_0}\rangle} \\
&= \bar\varphi^*(t_>;t_0)\,\bar\varphi(t_<;t_0)\,\frac{\langle 0_{t_0}|\,\bar a_0^\dagger\,\bar a_0\,|0_{t_0}\rangle}{\langle 0_{t_0}|0_{t_0}\rangle} \\
&= \frac{\mathrm{i}}{W_\rho[\bar\varphi^*(t;t_0),\,\bar\varphi(t;t_0)](T_s)}\,\bar\varphi^*(t_>;t_0)\,\bar\varphi(t_<;t_0)\,. \qquad (2.71)
\end{aligned}
$$

On the other hand, the last expression of (2.67) is obtained by using the following identities which are direct consequences of Eqs. (2.53), (2.56) and (2.62):

$$
\begin{aligned}
a_1 &= \bar\alpha(t_1;t_0)\,a_0 - \frac{\bar\beta(t_1;t_0)}{\alpha(t_1;t_0)}\,\big(\bar a_1 + \beta(t_1;t_0)\,a_0\big) \\
&= \frac{1}{\alpha(t_1;t_0)}\big(a_0 - \bar\beta(t_1;t_0)\,\bar a_1\big)\,, \qquad (2.72)
\end{aligned}
$$

$$
\begin{aligned}
\frac{\overline{\langle 0_{t_1}|}\,a_1\,\bar a_0\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle} &= \frac{1}{\alpha(t_1;t_0)}\,\frac{\overline{\langle 0_{t_1}|}\big(a_0 - \bar\beta(t_1;t_0)\,\bar a_1\big)\,\bar a_0\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle} = \frac{1}{\alpha(t_1;t_0)} \\
&= \frac{\mathrm{i}}{W_\rho[\varphi(t;t_1),\bar\varphi(t;t_0)]} \quad (t\text{: arbitrary})\,. \qquad (2.73)
\end{aligned}
$$

We then have

$$
\begin{aligned}
G_{10}(t,t';t_1,t_0) &= \frac{\overline{\langle 0_{t_1}|}\,\big(\varphi(t_>;t_1)\,a_1 + \bar\varphi(t_>;t_1)\,\bar a_1\big)\big(\varphi(t_<;t_0)\,a_0 + \bar\varphi(t_<;t_0)\,\bar a_0\big)|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle} \\
&= \varphi(t_>;t_1)\,\bar\varphi(t_<;t_0)\,\frac{\overline{\langle 0_{t_1}|}\,a_1\,\bar a_0\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle} \\
&= \frac{\mathrm{i}}{W_\rho[\varphi(t;t_1),\bar\varphi(t;t_0)]}\,\varphi(t_>;t_1)\,\bar\varphi(t_<;t_0) \quad (t\text{: arbitrary})\,. \qquad (2.74)
\end{aligned}
$$

When $\rho(t)$ and $\omega(t)$ are asymptotically constant in the remote past [i.e. $\rho(t) \sim \rho_{\rm in}$ and $\omega(t) \sim \omega_{\rm in}$ as $t \to t_i$], we can choose a pair of independent solutions $\{f(t), g(t)\}$ as those which behave as

$$
f(t) \sim e^{-\mathrm{i}\omega_{\rm in}t}\,, \qquad g(t) \sim e^{+\mathrm{i}\omega_{\rm in}t} \qquad (t \sim t_i)\,. \qquad (2.75)
$$

If we choose such basis, we then have

$$
u_0 \sim 0\,, \quad \bar u_0 \sim -2\mathrm{i}\rho_{\rm in}\,\omega_{\rm in}\,e^{-\mathrm{i}\omega_{\rm in}\,t_0}\,, \quad v_0 \sim 2\mathrm{i}\rho_{\rm in}\,\omega_{\rm in}\,e^{\mathrm{i}\omega_{\rm in}\,t_0}\,, \quad \bar v_0 \sim 0 \quad (t_0 \sim t_i)\,, \qquad (2.76)
$$

and from Eqs. (2.59) and (2.60) the wave functions at the remote past are found to behave as

$$
\varphi(t;t_0) \sim \frac{1}{\sqrt{2\rho_{\rm in}\,\omega_{\rm in}}}\,e^{-\mathrm{i}\omega_{\rm in}\,(t-t_0)}\,, \quad \bar\varphi(t;t_0) \sim \frac{1}{\sqrt{2\rho_{\rm in}\,\omega_{\rm in}}}\,e^{+\mathrm{i}\omega_{\rm in}\,(t-t_0)} \quad (t \sim t_i;\, t_0 \sim t_i)\,. \quad (2.77)
$$

A conclusion of the same kind can be obtained for the wave functions $\big(\varphi(t;t_1),\,\bar{\varphi}(t;t_1)\big)$ if $\rho(t)$ and $\omega(t)$ are asymptotically constant at the remote future. This behavior will be directly seen in concrete examples given in section 3 and in Appendix B.

3. Simple example: scalar field in Minkowski space

In this section, to demonstrate how the prescription of the previous section works, we consider a free real scalar field living in Minkowski space with the metric

$$\mathrm{d}s^2 = -\,\mathrm{d}t^2 + \mathrm{d}\boldsymbol{x}^2\,. \tag{3.1}$$

Another well studied example is investigated within our framework in Appendix B.

3.1. Setup

In order to clarify the structure of mode functions, we first assume that the spatial part is a $(d-1)$-dimensional torus of radius $L/2\pi$, which we will take infinite afterwards. The wave vectors $\boldsymbol{k}$ then take the following values:

$$\boldsymbol{k} = \frac{2\pi}{L}\,\boldsymbol{n} \quad \big(\boldsymbol{n}\in\mathbb{Z}^{d-1}\big)\,. \tag{3.2}$$

For $\boldsymbol{k} = \big(k_1, k_2, \ldots, k_{d-1}\big)$, we write $\boldsymbol{k} > 0$ (or $\boldsymbol{k} < 0$) if the first nonvanishing element in the sequence $\{k_1,\, k_2,\, \cdots\}$ is positive (or negative). Note that $\boldsymbol{k} < 0$ is equivalent to $-\boldsymbol{k} > 0$.

We introduce a complete set of (real-valued) eigenfunctions $\big\{Y_{\boldsymbol{k},\,a}(\boldsymbol{x})\big\}$ of the spatial Laplacian $\Delta_{d-1} = \sum_{i=1}^{d-1}\partial_i^2$ as

$$\boldsymbol{k} = 0: \quad Y_{\boldsymbol{k}=0,\,a=1} \equiv \frac{1}{\sqrt{V}} \quad \big(V \equiv L^{d-1}\big)\,, \tag{3.3}$$

$$\boldsymbol{k} > 0: \quad Y_{\boldsymbol{k},\,a=1}(\boldsymbol{x}) = \sqrt{\frac{2}{V}}\,\cos\boldsymbol{k}\cdot\boldsymbol{x}\,, \quad Y_{\boldsymbol{k},\,a=2}(\boldsymbol{x}) = \sqrt{\frac{2}{V}}\,\sin\boldsymbol{k}\cdot\boldsymbol{x}\,. \tag{3.4}$$

They satisfy the orthonormal relations,

$$\int \mathrm{d}^{d-1}\boldsymbol{x}\, Y_{\boldsymbol{k},\,a}(\boldsymbol{x})\, Y_{\boldsymbol{k}',\,a'}(\boldsymbol{x}) = \delta_{\boldsymbol{k},\boldsymbol{k}'}\,\delta_{a,a'}\,, \tag{3.5}$$

and we expand a scalar field $\phi(x)$ as

$$\phi(x) = \phi(t,\boldsymbol{x}) = \sum_{\boldsymbol{k}\ge 0}\sum_{a}\, \phi_{\boldsymbol{k},\,a}(t)\, Y_{\boldsymbol{k},\,a}(\boldsymbol{x})\,. \tag{3.6}$$

The action then becomes

$$\begin{aligned} S[\phi(x)] &= \int \mathrm{d}^d x \Big[-\frac{1}{2}\,\partial^\mu \phi(x)\,\partial_\mu \phi(x) - \frac{m^2 - \mathrm{i}\varepsilon}{2}\,\phi^2 \Big] \\ &= \sum_{\boldsymbol{k}\geq 0}\sum_a \int \mathrm{d}t \Big[\frac{1}{2}\,\dot{\phi}^2_{\boldsymbol{k},a}(t) - \frac{\omega^2_{k,\varepsilon}}{2}\,\phi^2_{\boldsymbol{k},a}(t) \Big], \end{aligned} \tag{3.7}$$

where

$$\omega_{k,\varepsilon} \equiv \sqrt{k^2 + m^2 - \mathrm{i}\varepsilon} = \omega_k - \mathrm{i}\varepsilon \quad \Big(k \equiv |\boldsymbol{k}| = \Big[\sum_{i=1}^{d-1} k_i^2\Big]^{1/2} \Big). \tag{3.8}$$

We thus have the following correspondence with the ingredients of the previous section:

$$q(t) = \phi_{\boldsymbol{k},a}(t)\,, \quad \rho(t) = 1\,, \quad \omega(t) = \omega_{k,\varepsilon} \ \ (= \text{constant in } t)\,. \tag{3.9}$$

3.2. Propagator for each mode

The equation of motion is given by $\ddot{q} + \omega^2 q = 0$, and we choose a pair of independent solutions as

$$f(t) = e^{-\mathrm{i}\omega t}\,, \quad g(t) = e^{\mathrm{i}\omega t}\,. \tag{3.10}$$

Their Wronskian is given by $W_\rho[f,g] = W[f,g] = 2\,\mathrm{i}\omega$, which is constant in t.

The functions $u(t)$ and $v(t)$ are easily found to be

$$\left\{ \begin{array}{c} u(t) \\ \bar{u}(t) \end{array} \right\} = \rho\,\big[\,\dot{f}(t) \pm \mathrm{i}\omega\, f(t)\,\big] = \left\{ \begin{array}{l} 0 \\ -2\,\mathrm{i}\,\omega\, e^{-\mathrm{i}\omega t} \end{array} \right., \tag{3.11}$$

$$\left\{ \begin{array}{c} v(t) \\ \bar{v}(t) \end{array} \right\} = \rho\,\big[\,\dot{g}(t) \pm \mathrm{i}\omega\, g(t)\,\big] = \left\{ \begin{array}{l} 2\,\mathrm{i}\,\omega\, e^{\mathrm{i}\omega t} \\ 0 \end{array} \right., \tag{3.12}$$

and using (2.59) and (2.60) we obtain the wave functions as

$$\varphi(t;t_0) = \frac{1}{W_\rho\,\sqrt{2\rho_0\,\omega_0}}\big[v_0\, f(t) - u_0\, g(t)\big] = \frac{1}{\sqrt{2\omega}}\,e^{-\mathrm{i}\omega\,(t-t_0)}\,, \tag{3.13}$$

$$\bar{\varphi}(t;t_0) = \frac{1}{W_\rho\,\sqrt{2\rho_0\,\omega_0}}\big[\bar{v}_0\, f(t) - \bar{u}_0\, g(t)\big] = \frac{1}{\sqrt{2\omega}}\,e^{\mathrm{i}\omega\,(t-t_0)}\,, \tag{3.14}$$

$$\varphi(t;t_1) = \frac{1}{W_\rho\,\sqrt{2\rho_1\,\omega_1}}\big[v_1\, f(t) - u_1\, g(t)\big] = \frac{1}{\sqrt{2\omega}}\,e^{-\mathrm{i}\omega\,(t-t_1)}\,, \tag{3.15}$$

$$\bar{\varphi}(t;t_1) = \frac{1}{W_\rho\,\sqrt{2\rho_1\,\omega_1}}\big[\bar{v}_1\, f(t) - \bar{u}_1\, g(t)\big] = \frac{1}{\sqrt{2\omega}}\,e^{\mathrm{i}\omega\,(t-t_1)}\,. \tag{3.16}$$

The Bogoliubov coefficients then can be calculated by using (2.62) as

$$\alpha(t_1;t_0)=e^{\mathrm{i}\omega\,(t_1-t_0)}\,,\quad \beta(t_1;t_0)=0\,,\quad \bar{\alpha}(t_1;t_0)=e^{-\mathrm{i}\omega\,(t_1-t_0)}\,,\quad \bar{\beta}(t_1;t_0)=0\,, \tag{3.17}$$

which indicates that the vacuum at a later time, $\overline{|0_{t_1}\rangle}$, coincides with the vacuum at an earlier time, $|0_{t_0}\rangle$, up to a phase. The two-point functions take the forms

$$\begin{aligned} G_{\boldsymbol{k},\,00}(t,t';t_0,t_0) &= \frac{\langle 0_{t_0}|\,\phi^\dagger_{\boldsymbol{k},\,a}(t_>)\,\phi_{\boldsymbol{k},\,a}(t_<)\,|0_{t_0}\rangle}{\langle 0_{t_0}|0_{t_0}\rangle} = \frac{\mathrm{i}\bar{\varphi}^*(t_>;t_0)\,\bar{\varphi}(t_<;t_0)}{W_\rho[\bar{\varphi}^*(t;t_0)\,\bar{\varphi}(t;t_0)](T_s)} \\ &= \frac{1}{2\,\mathrm{Re}\,\omega_{k,\,\varepsilon}\,e^{-2\varepsilon\,T_s}}\,e^{-\mathrm{i}\omega^*_{k,\,\varepsilon}\,t_>}\,e^{\mathrm{i}\omega_{k,\,\varepsilon}\,t_<}\,, \end{aligned} \tag{3.18}$$

$$\begin{aligned} G_{\boldsymbol{k},\,10}(t,t';t_1,t_0) &= \frac{\overline{\langle 0_{t_1}|}\,\mathrm{T}\,\phi_{\boldsymbol{k},\,a}(t)\,\phi_{\boldsymbol{k},\,a}(t')\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle} = \frac{1}{\alpha(t_1;t_0)}\,\varphi(t_>,t_1)\,\bar{\varphi}(t_<;t_0) \\ &= \frac{1}{2\,\omega_{k,\,\varepsilon}}\,e^{-\mathrm{i}\omega_{k,\,\varepsilon}\,(t_>-t_<)}\,, \end{aligned} \tag{3.19}$$

where we have restored the subscript of ω , $\omega_{k,\,\varepsilon}=\omega_k-\mathrm{i}\varepsilon$. Note that the dependence on t_0 and t_1 totally disappear in $G_{\boldsymbol{k},\,IJ}(t,t';t_I,t_J)$ for Minkowski space, and thus we need not to take the limit $t_0\to-\infty$, $t_1\to+\infty$ to obtain the in-in and in-out propagators. We find that the in-in and in-out propagators coincide after we take the limit $\varepsilon\to 0$,

$$G^{\mathrm{in/in}}_{\boldsymbol{k}}(t,t')=G^{\mathrm{in/out}}_{\boldsymbol{k}}(t,t')\,, \tag{3.20}$$

which we denote in the following discussions by

$$G_{\boldsymbol{k}}(t,t')=\frac{1}{2\,\omega_k}\,e^{-\mathrm{i}\omega_k\,(t_>-t_<)}\,\Big(=G_{-\boldsymbol{k}}(t,t')\Big)\,. \tag{3.21}$$

3.3. Propagator in spacetime

Once the propagators are obtained for each mode, the propagators in spacetime can be obtained by summing them over the modes $(\boldsymbol{k},a)$. The manipulation is known very well for Minkowski space, but we here review it briefly for later reference.

The in-in or in-out propagator is given by the following summation ($t_i=-\infty$, $t_f=\infty$):

$$G^{\mathrm{in/in}}(x,x')=\sum_{\boldsymbol{k},\,\boldsymbol{k}'\geq 0}\sum_{a,\,a'}\frac{\langle 0_{t_0}|\,\phi^\dagger_{\boldsymbol{k},\,a}(t_>)\,\phi_{\boldsymbol{k}',\,a'}(t_<)\,|0_{t_0}\rangle}{\langle 0_{t_0}|0_{t_0}\rangle}\Bigg|_{t_0\to t_i}\,Y_{\boldsymbol{k},\,a}(\boldsymbol{x})\,Y_{\boldsymbol{k}',\,a'}(\boldsymbol{x}')\,, \tag{3.22}$$

$$G^{\mathrm{in/out}}(x,x')=\sum_{\boldsymbol{k},\,\boldsymbol{k}'\geq 0}\sum_{a,\,a'}\frac{\overline{\langle 0_{t_1}|}\,\mathrm{T}\,\phi_{\boldsymbol{k},\,a}(t)\,\phi_{\boldsymbol{k}',\,a'}(t')\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle}\Bigg|_{\substack{t_0\to t_i\\ t_1\to t_f}}\,Y_{\boldsymbol{k},\,a}(\boldsymbol{x})\,Y_{\boldsymbol{k}',\,a'}(\boldsymbol{x}')\,. \tag{3.23}$$

Since the two-point functions are diagonalized with respect to the modes:

$$\left.\frac{\langle 0_{t_0}|\,\phi^\dagger_{\boldsymbol{k},\,a}(t_>)\,\phi_{\boldsymbol{k}',\,a'}(t_<)\,|0_{t_0}\rangle}{\langle 0_{t_0}|0_{t_0}\rangle}\right|_{t_0\to t_i} = G_{\boldsymbol{k}}(t,t')\,\delta_{\boldsymbol{k},\boldsymbol{k}'}\,\delta_{a,a'}\,, \tag{3.24}$$

$$\left.\frac{\overline{\langle 0_{t_1}|}\,\mathrm{T}\,\phi_{\boldsymbol{k},\,a}(t)\,\phi_{\boldsymbol{k}',\,a'}(t')\,|0_{t_0}\rangle}{\overline{\langle 0_{t_1}|}0_{t_0}\rangle}\right|_{\substack{t_0\to t_i\\ t_1\to t_f}} = G_{\boldsymbol{k}}(t,t')\,\delta_{\boldsymbol{k},\boldsymbol{k}'}\,\delta_{a,a'}\,, \tag{3.25}$$

we have

$$G^{\left\{\substack{\text{in/in}\\ \text{in/out}}\right\}}(x,x') = \sum_{\boldsymbol{k}\geq 0} G_{\boldsymbol{k}}(t,t') \sum_a Y_{\boldsymbol{k},a}(\boldsymbol{x})\,Y_{\boldsymbol{k},a}(\boldsymbol{x}') \equiv \sum_{\boldsymbol{k}\geq 0} G_{\boldsymbol{k}}(t,t')\,R_{\boldsymbol{k}}(\boldsymbol{x},\boldsymbol{x}')\,. \tag{3.26}$$

Here, $R_{\boldsymbol{k}}(\boldsymbol{x},\boldsymbol{x}') \equiv \sum_a Y_{\boldsymbol{k},a}(\boldsymbol{x})\,Y_{\boldsymbol{k},a}(\boldsymbol{x}')$ are easily calculated as

$$R_{\boldsymbol{k}=0}(\boldsymbol{x},\boldsymbol{x}') = \frac{1}{V}\,, \quad R_{\boldsymbol{k}>0}(\boldsymbol{x},\boldsymbol{x}') = \frac{2}{V}\,\cos\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{x})\,, \tag{3.27}$$

and thus we have

$$\begin{aligned} G(x,x') &= \frac{1}{V}\left[G_{\boldsymbol{k}=0}(t,t') + 2\sum_{\boldsymbol{k}>0} G_{\boldsymbol{k}}(t,t')\,\cos\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{x}')\right] \\ &= \frac{1}{V}\sum_{\boldsymbol{k}} G_{\boldsymbol{k}}(t,t')\,\cos\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{x}') \\ &= \int\frac{\mathrm{d}^{d-1}\boldsymbol{k}}{(2\pi)^{d-1}}\,G_{\boldsymbol{k}}(t,t')\,\cos\boldsymbol{k}\cdot(\boldsymbol{x}-\boldsymbol{x}')\,, \end{aligned} \tag{3.28}$$

where we have taken the limit $L\to\infty$ in the last equality.

The integration (3.28) can be performed easily (see Appendix C), and we obtain

$$\begin{aligned} G(x,x') &= \frac{1}{(2\pi)^{\frac{d-1}{2}}\,|\boldsymbol{x}-\boldsymbol{x}'|^{\frac{d-3}{2}}}\int_0^\infty \mathrm{d}k\,k^{\frac{d-1}{2}}\,G_{\boldsymbol{k}}(t,t')\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big) \\ &= \frac{m^{(d-2)/2}}{(2\pi)^{d/2}\,\big(\sigma+\mathrm{i}0\big)^{(d-2)/4}}\,K_{\frac{d-2}{2}}\big(m\sqrt{\sigma+\mathrm{i}0}\big) \end{aligned} \tag{3.29}$$

with $\sigma\equiv(x-x')^2$. Here, $J_\nu(z)$ is the Bessel function, and $K_\nu(z)$ is the modified Bessel function of the second kind.

4. Scalar field in de Sitter space

4.1. Geometry and definitions

We first recall the geometry of de Sitter space and collect the notation and definitions.

d-dimensional de Sitter space dS_d has the topology $\mathbb{R} \times S^{d-1}$ and is defined as a hyperboloid

$$\begin{aligned} &\eta_{MN}\, X^M X^N = \ell^2 \qquad (M, N, \cdots = 0, \ldots, d)\,, \\ &\bigl(\eta_{MN}\bigr) = \mathrm{diag}(-1, 1, \ldots, 1)\,, \end{aligned} \tag{4.1}$$

in $(d+1)$-dimensional Minkowski space with the metric

$$\mathrm{d}s^2 = \eta_{MN}\, \mathrm{d}X^M \mathrm{d}X^N\,. \tag{4.2}$$

ℓ is called the de Sitter radius. The constant Ricci scalar curvature is then given by $R = d\,(d-1)/\ell^2$.

There are several well-known coordinate patches which cover the whole or just a part of de Sitter space. Among them, we consider only the global patch and the Poincaré (or planer) patch, which we will briefly review below.

global patch: This coordinate patch covers the whole region of de Sitter space. The embedding of dS_d is given by the following functions:

$$X^0(\tau, \boldsymbol{\Omega}) = \ell\, \sinh\tau\,, \qquad X^I(\tau, \boldsymbol{\Omega}) = \ell\, \cosh\tau\, \Omega^I \quad (I = 1, \ldots, d) \tag{4.3}$$

with $\boldsymbol{\Omega}\cdot\boldsymbol{\Omega} = 1$. Here, τ runs over the range $-\infty < \tau < \infty$ and $\boldsymbol{\Omega}$ is a unit vector in $\mathbb{R}^d$ spanning a $(d-1)$-dimensional sphere. With the coordinates $(\tau, \boldsymbol{\Omega})$ the metric has the form

$$\mathrm{d}s^2 = \ell^2\bigl[-\mathrm{d}\tau^2 + \cosh^2\tau\, \mathrm{d}\Omega_{d-1}^2\bigr] = \ell^2\bigl[-\bigl(1-t^2\bigr)^{-2}\mathrm{d}t^2 + \bigl(1-t^2\bigr)^{-1}\mathrm{d}\Omega_{d-1}^2\bigr]\,. \tag{4.4}$$

In the last equality, we have introduced another temporal coordinate t as

$$t \equiv \tanh\tau \qquad (-1 < t < 1)\,. \tag{4.5}$$

Poincaré patch: This coordinate patch covers only half of de Sitter space. The embedding is given by the following functions with $\eta < 0$ and $\boldsymbol{x} \in \mathbb{R}^{d-1}$:

$$X^0(\eta, \boldsymbol{x}) = \frac{\ell^2 - \eta^2 + |\boldsymbol{x}|^2}{-2\eta}\,, \quad X^i(\eta, \boldsymbol{x}) = \ell\, \frac{x^i}{-\eta}\,, \quad X^d(\eta, \boldsymbol{x}) = \frac{\ell^2 + \eta^2 - |\boldsymbol{x}|^2}{-2\eta}\,, \tag{4.6}$$

where the spatial norm is defined by $|\boldsymbol{x}| \equiv \sqrt{\boldsymbol{x}\cdot\boldsymbol{x}} \equiv \sqrt{\delta_{ij}\, x^i\, x^j}$. Note that this patch only covers the region $X^0 + X^d = \ell^2/(-\eta) > 0$. In these coordinates, the metric takes the form

$$\mathrm{d}s^2 = \ell^2\, \frac{-\,\mathrm{d}\eta^2 + \mathrm{d}\boldsymbol{x}\cdot\mathrm{d}\boldsymbol{x}}{\eta^2}\,. \tag{4.7}$$

We define the de Sitter invariant quantity

$$Z(x,x') \equiv \ell^{-2}\,\eta_{MN}\,X^M(x)\,X^N(x')\,, \tag{4.8}$$

which is related to the geodesic distance $d(x,x')$ between two points x and x' via the relation

$$Z(x,x') = \cos\Bigl(\frac{d(x,x')}{\ell}\Bigr)\,. \tag{4.9}$$

It takes the following values depending on the positional relation between x and x' :

$$Z(x,x')\begin{cases} >1 & (\text{for } x \text{ and } x' \text{ timelikely separated}) \\ =1 & (\text{for } x \text{ and } x' \text{ lightlikely separated}) \\ <1 & (\text{for } x \text{ and } x' \text{ spacelikely separated}) \end{cases}\,, \tag{4.10}$$

as can be seen from the identity

$$Z(x,x') = 1 - \frac{1}{2\,\ell^2}\,\eta_{MN}\,\bigl(X^M(x)-X^M(x')\bigr)\bigl(X^N(x)-X^N(x')\bigr)\,. \tag{4.11}$$

In the global and the Poincaré coordinates, $Z(x,x')$ is written in the following form:

$$Z(x,x') = \begin{cases} -\sinh\tau\ \sinh\tau' + \cosh\tau\ \cosh\tau'\,\boldsymbol{\Omega}\cdot\boldsymbol{\Omega}' & (\text{global coords.}) \\ \dfrac{\eta^2+\eta'^2-|\boldsymbol{x}-\boldsymbol{x}'|^2}{2\eta\,\eta'} & (\text{Poincaré coords.}) \end{cases}\,. \tag{4.12}$$

Any de Sitter invariant quantity depending only on the geodesic distance $d(x,x')$ (such as Green's function of a de Sitter invariant vacuum) should be a function of $Z(x,x')$, and actually all the propagators considered in this paper can be written as functions of $Z(x,x')$. In what follows (except in subsec. 6.3), we set $\ell=1$.

4.2. Scalar field in the Poincaré patch

We first consider a free real scalar field in the Poincaré patch. The action takes the form[10]

$$S_\varepsilon[\phi(x)] = -\frac{1}{2}\int \mathrm{d}\eta \int \mathrm{d}^{d-1}\boldsymbol{x}\,\sqrt{-g}\,\bigl(g^{\mu\nu}\,\partial_\mu\phi\,\partial_\nu\phi + (m^2-\mathrm{i}\varepsilon)\,\phi^2\bigr)\,. \tag{4.13}$$

Using the same eigenfunctions $\{Y_{\boldsymbol{k},a}(\boldsymbol{x})\}$ as those given in section 3 [Eqs. (3.3), (3.4)], we expand a scalar field $\phi(x)$ as

$$\phi(x) = \phi(\eta,\boldsymbol{x}) = \sum_{\boldsymbol{k}\geq 0}\sum_a \phi_{\boldsymbol{k},a}(\eta)\,Y_{\boldsymbol{k},a}(\boldsymbol{x})\,. \tag{4.14}$$

---

[10] In this paper, we put a possible curvature-coupling term, $(\xi/2)\,R\,\phi^2 = \bigl(d(d-1)\xi/2\bigr)\,\phi^2$, into the mass term, $(m^2/2)\,\phi^2$.

The functions defined in section 2 then take the following form (see (2.8)):

$$q(\eta)=\phi_{\boldsymbol{k},a}(\eta)\,,\quad \rho(\eta)=(-\eta)^{-(d-2)}\,, \tag{4.15}$$

$$\omega(\eta)=\sqrt{(m^2-\mathrm{i}\varepsilon)\,(-\eta)^{-2}+k^2}\quad (k\equiv|\boldsymbol{k}|)\,, \tag{4.16}$$

from which we introduce[11]

$$\omega_0\equiv\omega(\eta_0)\overset{\eta_0\to-\infty}{\longrightarrow}k\,,\qquad \omega_1\equiv\omega(\eta_1)\overset{\eta_1\to-0}{\longrightarrow}\frac{\bar m_1}{-\eta_1}\,, \tag{4.17}$$

$$\bar m_1\equiv\begin{cases} m-\mathrm{i}\varepsilon & (\text{for } m>0)\\ -\eta_1\,k-\mathrm{i}\varepsilon & (\text{for } m=0)\end{cases}\,. \tag{4.18}$$

4.2.1. Propagators for each mode in the Poincaré patch

The equation of motion (2.35) takes the form

$$\eta^2\,\ddot q(\eta)-(d-2)\,\dot q(\eta)+\bigl(k^2\eta^2+m^2-\mathrm{i}\varepsilon\bigr)\,q(\eta)=0\,, \tag{4.19}$$

and we choose a set of independent solutions as

$$f(\eta)=(-\eta)^{\frac{d-1}{2}}\,J_{\nu_\varepsilon}(-k\,\eta)\,,\qquad g(\eta)=(-\eta)^{\frac{d-1}{2}}\,N_{\nu_\varepsilon}(-k\,\eta) \tag{4.20}$$

with

$$\nu_\varepsilon\equiv\begin{cases}\sqrt{\bigl(\frac{d-1}{2}\bigr)^2-(m^2-\mathrm{i}\varepsilon)}=\nu+\mathrm{i}\varepsilon & \bigl(m<\frac{d-1}{2}\bigr)\\ \mathrm{i}\sqrt{m^2-\mathrm{i}\varepsilon-\bigl(\frac{d-1}{2}\bigr)^2}=\mathrm{i}\,\mu+\varepsilon & \bigl(m\ge\frac{d-1}{2}\bigr)\end{cases}$$
$$\left[\nu\equiv\sqrt{\Bigl(\frac{d-1}{2}\Bigr)^2-m^2}\,,\qquad \mu\equiv\sqrt{m^2-\Bigl(\frac{d-1}{2}\Bigr)^2}\right]. \tag{4.21}$$

Here, $N_\nu(x)$ is the Neumann function. Note that $\mathrm{Re}\,\nu_\varepsilon>0$ for any positive value of m. The Wronskian is given by $W[f,g](\eta)=-(2/\pi)\,(-\eta)^{d-2}$, and thus

$$W_\rho[f,g]=\rho(t)\,W[f,g](t)=-\frac{2}{\pi}\,. \tag{4.22}$$

The functions $u(\eta)$ and $v(\eta)$ of (2.49) have the form

$$\left\{\begin{matrix}u(\eta)\\ \bar u(\eta)\end{matrix}\right\}=-(-\eta)^{-\frac{d-1}{2}}\Bigl[\Bigl(\frac{d-1}{2}+\nu_\varepsilon\pm\mathrm{i}\omega\,\eta\Bigr)J_{\nu_\varepsilon}(-k\,\eta)+k\,\eta\,J_{1+\nu_\varepsilon}(-k\,\eta)\Bigr]\,, \tag{4.23}$$

$$\left\{\begin{matrix}v(\eta)\\ \bar v(\eta)\end{matrix}\right\}=-(-\eta)^{-\frac{d-1}{2}}\Bigl[\Bigl(\frac{d-1}{2}+\nu_\varepsilon\pm\mathrm{i}\omega\,\eta\Bigr)N_{\nu_\varepsilon}(-k\,\eta)+k\,\eta\,N_{1+\nu_\varepsilon}(-k\,\eta)\Bigr]\,, \tag{4.24}$$

---

[11] Our prescription of section 2 cannot be applied directly to the exactly massless case, where $\omega(\eta)=0$ for $k=0$. We actually define the massless theory as the $m\to0$ limit of a massive theory, and the expression for $\bar m_1$ in the massless case is presented just for reference.

where we have used the formulae

$$z\,\frac{\partial J_\nu(z)}{\partial z} = \nu\,J_\nu(z) - z\,J_{\nu+1}(z)\,, \qquad z\,\frac{\partial N_\nu(z)}{\partial z} = \nu\,N_\nu(z) - z\,N_{\nu+1}(z)\,. \tag{4.25}$$

The wave functions are then given by

$$\varphi(\eta;\eta_I) = -\frac{\pi}{2\sqrt{2\omega_I}}\,\frac{(-\eta)^{\frac{d-1}{2}}}{(-\eta_I)^{-\frac{d-2}{2}}}\,\bigl(v_I\,J_{\nu_\varepsilon}(-k\,\eta) - u_I\,N_{\nu_\varepsilon}(-k\,\eta)\bigr)\,, \tag{4.26}$$

$$\overline{\varphi}(\eta;\eta_I) = -\frac{\pi}{2\sqrt{2\omega_I}}\,\frac{(-\eta)^{\frac{d-1}{2}}}{(-\eta_I)^{-\frac{d-2}{2}}}\,\bigl(\bar{v}_I\,J_{\nu_\varepsilon}(-k\,\eta) - \bar{u}_I\,N_{\nu_\varepsilon}(-k\,\eta)\bigr)\,, \tag{4.27}$$

with which we obtain an exact expression of the propagators for a finite time interval:

$$\begin{aligned}
&G_{\boldsymbol{k},00}(\eta,\eta';\eta_0,\eta_0)\\
&= \begin{cases}
\dfrac{-\mathrm{i}\pi\,[(-\eta)(-\eta')]^{\frac{d-1}{2}}}{2(\bar{u}_0^*\,\bar{v}_0 - \bar{v}_0^*\,\bar{u}_0)} \\
\times\bigl[\bar{v}_0^*\,J_{\nu_{-\varepsilon}}(-k\,\eta') - \bar{u}_0^*\,N_{\nu_{-\varepsilon}}(-k\,\eta')\bigr]\,\bigl[\bar{v}_0\,J_{\nu_\varepsilon}(-k\,\eta') - \bar{u}_0\,N_{\nu_\varepsilon}(-k\,\eta')\bigr] & \Bigl(m < \dfrac{d-1}{2}\Bigr)\\
\dfrac{-\mathrm{i}\pi\,[(-\eta)(-\eta')]^{\frac{d-1}{2}}}{2\bigl[(\bar{u}_0^*\,\bar{v}_0 - \bar{v}_0^*\,\bar{u}_0)\cos(\pi\nu_{-\varepsilon}) + (|\bar{u}_0|^2 + |\bar{v}_0|^2)\sin(\pi\nu_{-\varepsilon})\bigr]} \\
\times\bigl[\bar{v}_0^*\,J_{-\nu_{-\varepsilon}}(-k\,\eta') - \bar{u}_0^*\,N_{-\nu_{-\varepsilon}}(-k\,\eta')\bigr]\,\bigl[\bar{v}_0\,J_{\nu_\varepsilon}(-k\,\eta') - \bar{u}_0\,N_{\nu_\varepsilon}(-k\,\eta')\bigr] & \Bigl(m \geq \dfrac{d-1}{2}\Bigr)
\end{cases},
\end{aligned} \tag{4.28}$$

$$\begin{aligned}
&G_{\boldsymbol{k},10}(\eta,\eta';\eta_1,\eta_0)\\
&= \frac{-\mathrm{i}\pi\,[(-\eta)(-\eta')]^{\frac{d-1}{2}}}{2(u_1\,\bar{v}_0 - v_1\,\bar{u}_0)}\,\bigl[v_1\,J_{\nu_\varepsilon}(-k\,\eta) - u_1\,N_{\nu_\varepsilon}(-k\,\eta)\bigr]\bigl[\bar{v}_0\,J_{\nu_\varepsilon}(-k\,\eta') - \bar{u}_0\,N_{\nu_\varepsilon}(-k\,\eta')\bigr]\,.
\end{aligned} \tag{4.29}$$

We now send η_0, η_1 to the boundary of the Poincaré patch; $\eta_0 \to -\infty$ and $\eta_1 \to 0$. The wave functions $\varphi(\eta;\eta_I)$ and $\bar{\varphi}(\eta;\eta_I)$ $(I = 0, 1)$ then behave as

$$\varphi(\eta;\eta_0) \sim \frac{\sqrt{\pi}}{2}\,e^{\mathrm{i}\left(k\,\eta_0 + \frac{\pi(2\nu_\varepsilon+1)}{4}\right)}\,(-\eta)^{\frac{d-1}{2}}\,H^{(1)}_{\nu_\varepsilon}(-k\,\eta)\,, \tag{4.30}$$

$$\bar{\varphi}(\eta;\eta_0) \sim \frac{\sqrt{\pi}}{2}\,e^{-\mathrm{i}\left(k\,\eta_0 + \frac{\pi(2\nu_\varepsilon+1)}{4}\right)}\,(-\eta)^{\frac{d-1}{2}}\,H^{(2)}_{\nu_\varepsilon}(-k\,\eta)\,, \tag{4.31}$$

$$\varphi(\eta;\eta_1) \sim -\frac{\Gamma(\nu_\varepsilon)\,(k/2)^{-\nu_\varepsilon}}{2\sqrt{2\bar{m}_1}}\,(-\eta_1)^{-\nu_\varepsilon}\,\Bigl(\frac{d-1}{2} - \nu_\varepsilon - \mathrm{i}\bar{m}_1\Bigr)\,(-\eta)^{\frac{d-1}{2}}\,J_{\nu_\varepsilon}(-k\,\eta)\,, \tag{4.32}$$

$$\bar{\varphi}(\eta;\eta_1) \sim -\frac{\Gamma(\nu_\varepsilon)\,(k/2)^{-\nu_\varepsilon}}{2\sqrt{2\bar{m}_1}}\,(-\eta_1)^{-\nu_\varepsilon}\,\Bigl(\frac{d-1}{2} - \nu_\varepsilon + \mathrm{i}\bar{m}_1\Bigr)\,(-\eta)^{\frac{d-1}{2}}\,J_{\nu_\varepsilon}(-k\,\eta)\,. \tag{4.33}$$

Here, $H^{(1)}_\nu(x)$ and $H^{(2)}_\nu(x)$ are the Hankel functions defined by $H^{(1,2)}_\nu(x) \equiv J_\nu(x) \pm \mathrm{i}N_\nu(x)$.

They can be obtained by using the asymptotic forms

$$\left\{\begin{array}{c} u_0 \\ \bar{u}_0 \end{array}\right\} \sim \pm\mathrm{i}\sqrt{\frac{2k}{\pi}}\,(-\eta_0)^{-\frac{d-2}{2}} e^{\pm\mathrm{i}\left(k\,\eta_0+\frac{\pi(2\nu_\varepsilon+1)}{4}\right)}\,, \tag{4.34}$$

$$\left\{\begin{array}{c} v_0 \\ \bar{v}_0 \end{array}\right\} \sim -\sqrt{\frac{2k}{\pi}}\,(-\eta_0)^{-\frac{d-2}{2}} e^{\pm\mathrm{i}\left(k\,\eta_0+\frac{\pi(2\nu_\varepsilon+1)}{4}\right)}\,, \tag{4.35}$$

$$\left\{\begin{array}{c} u_1 \\ \bar{u}_1 \end{array}\right\} \sim -\frac{(k/2)^{\nu_\varepsilon}}{\Gamma(1+\nu_\varepsilon)}(-\eta_1)^{-\frac{d-1}{2}+\nu_\varepsilon}\left(\frac{d-1}{2}+\nu_\varepsilon\mp\mathrm{i}\bar{m}_1\right), \tag{4.36}$$

$$\left\{\begin{array}{c} v_1 \\ \bar{v}_1 \end{array}\right\} \sim \frac{\Gamma(\nu_\varepsilon)\,(k/2)^{-\nu_\varepsilon}}{\pi}\,(-\eta_1)^{-\frac{d-1}{2}-\nu_\varepsilon}\left(\frac{d-1}{2}-\nu_\varepsilon\mp\mathrm{i}\bar{m}_1\right), \tag{4.37}$$

which can be easily derived by using the asymptotic forms of the Bessel functions:

$$J_{\nu_\varepsilon}(-k\,\eta) \overset{\eta\to 0}{\sim} \frac{1}{\Gamma(1+\nu_\varepsilon)}\left(-\frac{k\,\eta}{2}\right)^{\nu_\varepsilon}, \qquad N_{\nu_\varepsilon}(-k\,\eta) \overset{\eta\to 0}{\sim} -\frac{\Gamma(\nu_\varepsilon)}{\pi}\left(-\frac{k\,\eta}{2}\right)^{-\nu_\varepsilon}, \tag{4.38}$$

$$J_{\nu_\varepsilon}(-k\,\eta) \overset{\eta\to -\infty}{\sim} \sqrt{\frac{2}{\pi}}\,(-k\,\eta)^{-1/2}\cos\bigl[-k\,\eta-(\pi/4)(1+2\nu_\varepsilon)\bigr]\,, \tag{4.39}$$

$$N_{\nu_\varepsilon}(-k\,\eta) \overset{\eta\to -\infty}{\sim} \sqrt{\frac{2}{\pi}}\,(-k\,\eta)^{-1/2}\sin\bigl[-k\,\eta-(\pi/4)(1+2\nu_\varepsilon)\bigr]\,. \tag{4.40}$$

Note that the wave function $\varphi(\eta;\eta_0)$ converges to a finite function,

$$\varphi_{\rm in}(\eta) \equiv \frac{\sqrt{\pi}}{2}\,(-\eta)^{\frac{d-1}{2}}\,H^{(1)}_{\nu_\varepsilon}(-k\,\eta)\,, \tag{4.41}$$

in the limit $\eta_0 \to -\infty$ (up to an oscillatory phase), while $\varphi(\eta;\eta_1)$ is divergent in the limit $\eta_1 \to 0$. This difference can be attributed to the fact that the timelike vector ∂_η becomes asymptotically a Killing vector in the remote past, but does not in the remote future.[12] The finite asymptotic function $\varphi_{\rm in}(\eta)$ coincides with the positive-mode wave function associated with the Euclidean vacuum up to a phase.

The Bogoliubov coefficients have the asymptotic forms

$$\left\{\begin{array}{c} \alpha(\eta_1;\eta_0) \\ \beta(\eta_1;\eta_0) \end{array}\right\} \sim -\frac{\Gamma(\nu_\varepsilon)}{2\sqrt{2\pi}\,\bar{m}_1}\,(-k\,\eta_1/2)^{-\nu_\varepsilon}\left(\frac{d-1}{2}-\nu_\varepsilon-\mathrm{i}\bar{m}_1\right)e^{\mp\mathrm{i}\left(k\,\eta_0+\frac{\pi(2\nu_\varepsilon+1)}{4}\right)}\,, \tag{4.42}$$

where we have used the approximation $(-\eta_1)^{-\nu_\varepsilon} \gg (-\eta_1)^{\nu_\varepsilon}$, since ν_ε always has a positive real part. Note that the norm of the Bogoliubov coefficients $|\alpha(\eta_1;\eta_0)|$ and $|\beta(\eta_1;\eta_0)|$ diverges in the limit $\eta_1 \to 0$ although the relation (2.56) still holds. This is an evidence that the remote future cannot be treated adiabatically [7].

---

[12] In fact, the Lie derivative of $g_{\mu\nu}$ with respect to the vector $\xi = \partial_\eta$ is $\pounds_\xi\, g_{\mu\nu} \propto (-\eta)^{-3}$.

Substituting everything into Eqs. (2.66) and (2.67) and taking the limit $\eta_0 \to -\infty$ and $\eta_1 \to 0$, we obtain the propagators in the Poincaré patch,

$$G_{\boldsymbol{k}}^{\mathrm{in/in}}(\eta,\eta') = \frac{\pi}{4}\,[(-\eta)(-\eta')]^{\frac{d-1}{2}}\,H^{(1)}_{\nu_{-\varepsilon}}\big(-k\,\eta_>\big)\,H^{(2)}_{\nu_\varepsilon}\big(-k\,\eta_<\big)\,, \tag{4.43}$$

$$G_{\boldsymbol{k}}^{\mathrm{in/out}}(\eta,\eta') = \frac{\pi}{2}\,[(-\eta)(-\eta')]^{\frac{d-1}{2}}\,J_{\nu_\varepsilon}\big(-k\,\eta_>\big)\,H^{(2)}_{\nu_\varepsilon}\big(-k\,\eta_<\big)\,. \tag{4.44}$$

One should note that, in the limit $\eta_1 \to 0$, the divergence in $\varphi(\eta;\eta_1)$ is canceled out with that of $\alpha(\eta_1;\eta_0)$ and the in-out propagator is obtained with a finite value.

4.2.2. Propagators in the Poincaré patch

Since the eigenfunctions $\{Y_{\boldsymbol{k},a}(\boldsymbol{x})\}$ are the same as those given in section 3, the propagators in spacetime can be written in the form

$$\begin{aligned} G^{\left\{\substack{\mathrm{in/in}\\ \mathrm{in/out}}\right\}}(x,x') &= \sum_{\boldsymbol{k}\geq 0} G_{\boldsymbol{k}}^{\left\{\substack{\mathrm{in/in}\\ \mathrm{in/out}}\right\}}(\eta,\eta') \sum_a Y_{\boldsymbol{k},a}(\boldsymbol{x})\,Y_{\boldsymbol{k},a}(\boldsymbol{x}') \\ &= \frac{1}{(2\pi)^{\frac{d-1}{2}}\,|\boldsymbol{x}-\boldsymbol{x}'|^{\frac{d-3}{2}}} \int_0^\infty \mathrm{d}k\,k^{\frac{d-1}{2}}\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big)\,G_{\boldsymbol{k}}^{\left\{\substack{\mathrm{in/in}\\ \mathrm{in/out}}\right\}}(\eta,\eta')\,, \end{aligned} \tag{4.45}$$

as in the case of Minkowski space (see Appendix C).

For the in-in propagator, we have

$$\begin{aligned} &G^{\mathrm{in/in}}(x,x') \\ &= \frac{\pi}{4}\,\frac{[(-\eta)(-\eta')]^{\frac{d-1}{2}}}{(2\pi)^{\frac{d-1}{2}}\,|\boldsymbol{x}-\boldsymbol{x}'|^{\frac{d-3}{2}}} \int_0^\infty \mathrm{d}k\,k^{\frac{d-1}{2}}\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big)\,H^{(1)}_{\nu_{-\varepsilon}}(-k\,\eta_>)\,H^{(2)}_{\nu_\varepsilon}(-k\,\eta_<)\,. \end{aligned} \tag{4.46}$$

As is proved in Appendix D, this can be integrated to the form[13]

$$G^{\mathrm{in/in}}(x,x') = \frac{\Gamma\big(\frac{d-1}{2}+\nu\big)\,\Gamma\big(\frac{d-1}{2}-\nu\big)}{2\,(2\pi)^{d/2}}\,(u^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}^{-\frac{d-2}{2}}_{\nu-1/2}(u) \tag{4.47}$$

with $u \equiv -Z(x,x') + \mathrm{i}\,0$. This can be further rewritten as

$$\begin{aligned} G^{\mathrm{in/in}}(x,x') &= \frac{\Gamma\big(\frac{d-1}{2}+\nu\big)\,\Gamma\big(\frac{d-1}{2}-\nu\big)}{(4\pi)^{d/2}\,\Gamma\big(d/2\big)}\,F\Big(\frac{d-1}{2}+\nu\,,\,\frac{d-1}{2}-\nu\,;\,\frac{d}{2}\,;\,\frac{1-u}{2}\Big) \qquad (4.48) \\ &= \frac{\Gamma\big(\frac{d-1}{2}\big)}{4\pi^{(d-1)/2}\,\sin\big[\pi\big(\frac{d-1}{2}-\nu\big)\big]}\,C^{\frac{d-1}{2}}_{\nu-\frac{d-1}{2}}(u)\,, \qquad (4.49) \end{aligned}$$

---

[13] Here we have set $\varepsilon = 0$. $\mathcal{P}^\mu_\nu(z)$ and $\mathcal{Q}^\mu_\nu(z)$ denote the associated Legendre functions of the first and second kind that are defined on the complex z-plane other than the cut along the real axis to the left of the point $z = 1$. There are another type of associated Legendre functions that are defined on the interval $(-1,1)$, which we denote by $\mathsf{P}(x)$ and $\mathsf{Q}(x)$. See Appendix E for their definitions and several useful identities.

where Kummer's relation (B.29) has been used in the first equality and $C_\nu^\mu(x)$ is the Gegenbauer function. This propagator is the same as the well-known in-in propagator associated with the Euclidean vacuum.

The in-out propagator

$$\begin{aligned} &G^{\mathrm{in/out}}(x,x') \\ &= \frac{\pi}{2}\,\frac{[(-\eta)(-\eta')]^{\frac{d-1}{2}}}{(2\pi)^{\frac{d-1}{2}}\,|\boldsymbol{x}-\boldsymbol{x}'|^{\frac{d-3}{2}}}\int_0^\infty \mathrm{d}k\,k^{\frac{d-1}{2}}\,J_{\frac{d-3}{2}}\bigl(k|\boldsymbol{x}-\boldsymbol{x}'|\bigr)\,J_{\nu_\varepsilon}(-k\,\eta_>)\,H^{(2)}_{\nu_\varepsilon}(-k\,\eta_<) \end{aligned} \tag{4.50}$$

can be rewritten in a similar manner to the form

$$G^{\mathrm{in/out}}(x,x') = \frac{e^{-\mathrm{i}\pi(d-2)}}{(2\pi)^{d/2}}\,(u^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-1/2}(u) \tag{4.51}$$

with $u = Z(x,x') - \mathrm{i}0$. A proof is given also in Appendix D.

If we consider the massless limit where $\nu \to (d-1)/2$, the in-in propagator diverges, as was pointed out in [3]. In contrast, we find that the in-out propagator has a finite massless limit:

$$G^{\mathrm{in/out}}(x,x') = \frac{e^{-\mathrm{i}\pi(d-2)}}{(2\pi)^{d/2}}\,(u^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\frac{d-2}{2}}(u) \quad (\text{massless})\,. \tag{4.52}$$

4.3. Scalar field in the global patch

In the global patch, we choose a set of (real-valued) spherical harmonics $\{Y_{LM}(\boldsymbol{\Omega})\}$ as the complete eigenfunctions of the spatial Laplacian Δ_{d-1} on S^{d-1}. They satisfy $\Delta_{d-1}\,Y_{LM} = -L(L+d-2)\,Y_{LM}$ $(M=1,\ldots,N_L^{(d)})$, and the degeneracy $N_L^{(d)}$ is given by

$$N_L^{(d)} = \frac{(L+d-3)!}{(d-2)!\,L!}\,(2L+d-2) \tag{4.53}$$

with the exception of the case $d=2$ and $L=0$ where $N_0^{(2)} \equiv 1$. We choose them such that they are orthonormal:

$$\int \mathrm{d}\boldsymbol{\Omega}\,Y_{LM}(\boldsymbol{\Omega})\,Y_{L'M'}(\boldsymbol{\Omega}) = \delta_{LL'}\,\delta_{MM'}\,. \tag{4.54}$$

Then, by expanding $\phi(x)$ as

$$\phi(x) = \sum_{L,M}\phi_{LM}(t)\,Y_{LM}(\boldsymbol{\Omega})\,, \tag{4.55}$$

the mode function $q(t) \equiv \phi_{LM}(t)$ describes a harmonic oscillator with time-dependent mass and frequency of the following form [see (2.8)]:

$$\rho(t) = \big(1 - t^2\big)^{-\frac{d-3}{2}}\,, \tag{4.56}$$

$$\omega(t) = \big(1 - t^2\big)^{-1} \sqrt{L\,(L + d - 2)\,\big(1 - t^2\big) + m^2 - \mathrm{i}\varepsilon} \equiv \big(1 - t^2\big)^{-1}\,\bar{m}(t)\,. \tag{4.57}$$

4.3.1. Propagators for each mode in the global patch

We choose a pair of independent solutions as

$$f(t) = \big(1 - t^2\big)^{\frac{d-1}{4}}\,\mathsf{P}_k^{\nu_\varepsilon}(t)\,, \qquad g(t) = \big(1 - t^2\big)^{\frac{d-1}{4}}\,\mathsf{Q}_k^{\nu_\varepsilon}(t)\,, \tag{4.58}$$

where

$$k \equiv L + \frac{d-3}{2}\,, \tag{4.59}$$

and $\mathsf{P}_\nu^\mu(t)$ and $\mathsf{Q}_\nu^\mu(t)$ are the associated Legendre functions defined on the interval $(-1, 1)$ (see Appendix E). The weighted Wronskian (constant in t) then has the form

$$W_\rho[f, g] = \rho(t)\,W[f, g](t) = \frac{\Gamma(k + \nu_\varepsilon + 1)}{\Gamma(k - \nu_\varepsilon + 1)}\,. \tag{4.60}$$

The functions $u(t)$ and $v(t)$ defined in (2.49) are then given by

$$\left\{ \begin{array}{c} u(t) \\ \bar{u}(t) \end{array} \right\} = \big(1 - t^2\big)^{-\frac{d-1}{4}} \Big[\big(L\,t \pm \mathrm{i}\bar{m}(t)\big)\mathsf{P}_k^{\nu_\varepsilon}(t) - \big(k - \nu_\varepsilon + 1\big)\mathsf{P}_{k+1}^{\nu_\varepsilon}(t) \Big]\,, \tag{4.61}$$

$$\left\{ \begin{array}{c} v(t) \\ \bar{v}(t) \end{array} \right\} = \big(1 - t^2\big)^{-\frac{d-1}{4}} \Big[\big(L\,t \pm \mathrm{i}\bar{m}(t)\big)\mathsf{Q}_k^{\nu_\varepsilon}(t) - \big(k - \nu_\varepsilon + 1\big)\mathsf{Q}_{k+1}^{\nu_\varepsilon}(t) \Big]\,, \tag{4.62}$$

and the wave functions $(\varphi,\, \bar{\varphi})$ take the form

$$\varphi(t; t_I) = \frac{\Gamma(k - \nu_\varepsilon + 1)}{\sqrt{2\bar{m}_I}\,\Gamma(k + \nu_\varepsilon + 1)}\,(1 - t_I^2)^{\frac{d-1}{4}}\big(v_I\,f(t) - u_I\,g(t)\big)\,, \tag{4.63}$$

$$\bar{\varphi}(t; t_I) = \frac{\Gamma(k - \nu_\varepsilon + 1)}{\sqrt{2\bar{m}_I}\,\Gamma(k + \nu_\varepsilon + 1)}\,(1 - t_I^2)^{\frac{d-1}{4}}\big(\bar{v}_I\,f(t) - \bar{u}_I\,g(t)\big)\,. \tag{4.64}$$

We now send t_0, t_1 to the boundary of the global patch; $t_0 \to -1$ and $t_1 \to 1$. Using Eqs. (E.21) and (E.22), we see that $u_1 = u(t_1)$ and $v_1 = v(t_1)$ take the following asymptotic forms in the limit $t_1 \to 1$:

$$\left\{ \begin{array}{c} u_1 \\ \bar{u}_1 \end{array} \right\} \sim \frac{2^{\nu_\varepsilon}\,\sin(\pi\nu_\varepsilon)\,\Gamma(\nu_\varepsilon)}{\pi}\Big(-\frac{d-1}{2} + \nu_\varepsilon \pm \mathrm{i}\bar{m}_1\Big)\,\big(1 - t_1^2\big)^{-\frac{d-1}{4} - \frac{\nu_\varepsilon}{2}}\,, \tag{4.65}$$

$$\left\{ \begin{array}{c} v_1 \\ \bar{v}_1 \end{array} \right\} \sim 2^{\nu_\varepsilon - 1}\,\cos(\pi\nu_\varepsilon)\,\Gamma(\nu_\varepsilon)\,\Big(-\frac{d-1}{2} + \nu_\varepsilon \pm \mathrm{i}\bar{m}_1\Big)\,\big(1 - t_1^2\big)^{-\frac{d-1}{4} - \frac{\nu_\varepsilon}{2}}\,, \tag{4.66}$$

and thus we find that the wave functions behave as[14]

$$\left\{ \begin{array}{c} \varphi(t;t_1) \\ \bar{\varphi}(t;t_1) \end{array} \right\} \sim \left[\frac{2^{\nu_\varepsilon-1}\,\Gamma(\nu_\varepsilon)}{\sqrt{2\bar{m}_1}} \Big(-\frac{d-1}{2} + \nu_\varepsilon \pm \mathrm{i}\bar{m}_1 \Big) \big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}} \right] \big(1-t^2\big)^{\frac{d-1}{4}}\, \mathsf{P}_k^{-\nu_\varepsilon}(t)\,. \tag{4.67}$$

Here, we have used the fact that $\bar{m}_1 \sim m - \mathrm{i}\varepsilon$ in this region.

Similarly, $u_0 = u(t_0)$ and $v_0 = v(t_0)$ have the following asymptotic forms in the limit $t_0 \to -1$:

$$\left\{ \begin{array}{c} u_0 \\ \bar{u}_0 \end{array} \right\} \sim \left\{ \begin{array}{ll} \dfrac{2^{-\nu_\varepsilon}\,\cos(\pi k)\,\Gamma(k+\nu_\varepsilon+1)}{\Gamma(\nu_\varepsilon+1)\,\Gamma(k-\nu_\varepsilon+1)} \Big(-\dfrac{d-1}{2} + \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \Big(1-t_0^2\Big)^{-\frac{d-1}{4}+\frac{\nu_\varepsilon}{2}} & (d : \text{odd}) \\ -\dfrac{2^{\nu_\varepsilon}\,\sin(\pi k)\,\Gamma(\nu_\varepsilon)}{\pi} \Big(\dfrac{d-1}{2} - \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \big(1-t_0^2\big)^{-\frac{d-1}{4}-\frac{\nu_\varepsilon}{2}} & (d : \text{even}) \end{array} \right. , \tag{4.68}$$

$$\left\{ \begin{array}{c} v_0 \\ \bar{v}_0 \end{array} \right\} \sim \left\{ \begin{array}{ll} -\,2^{\nu_\varepsilon-1}\,\cos(\pi k)\,\Gamma(\nu_\varepsilon) \Big(\dfrac{d-1}{2} - \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \big(1-t_0^2\big)^{-\frac{d-1}{4}-\frac{\nu_\varepsilon}{2}} & (d : \text{odd}) \\ \dfrac{2^{-\nu_\varepsilon-1}\,\pi\,\Gamma(k+\nu_\varepsilon+1)}{\sin(\pi k)\,\Gamma(\nu_\varepsilon+1)\,\Gamma(k-\nu_\varepsilon+1)} \Big(\dfrac{d-1}{2} - \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \big(1-t_0^2\big)^{-\frac{d-1}{4}+\frac{\nu_\varepsilon}{2}} & (d : \text{even}) \end{array} \right. . \tag{4.69}$$

Here, we have used the relation $\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}} \gg \big(1-t_0^2\big)^{\frac{\nu_\varepsilon}{2}}$, which holds since $\mathrm{Re}\,\nu_\varepsilon > 0$. We thus find that the wave functions behave as

$$\left\{ \begin{array}{c} \varphi(t;t_0) \\ \bar{\varphi}(t;t_0) \end{array} \right\} \sim \left[-\frac{2^{\nu_\varepsilon-1}\,\cos(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\sqrt{2\bar{m}_0}\,\Gamma(k+\nu_\varepsilon+1)} \Big(\frac{d-1}{2} - \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}} \right] \times \big(1-t^2\big)^{\frac{d-1}{4}}\, \mathsf{P}_k^{\nu_\varepsilon}(t) \tag{4.70}$$

in odd dimensions, and

$$\left\{ \begin{array}{c} \varphi(t;t_0) \\ \bar{\varphi}(t;t_0) \end{array} \right\} \sim \left[\frac{2^{\nu_\varepsilon}\,\sin(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\pi\,\sqrt{2\bar{m}_0}\,\Gamma(k+\nu_\varepsilon+1)} \Big(\frac{d-1}{2} - \nu_\varepsilon \pm \mathrm{i}\bar{m}_0 \Big) \big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}} \right] \times \big(1-t^2\big)^{\frac{d-1}{4}}\, \mathsf{Q}_k^{\nu_\varepsilon}(t) \tag{4.71}$$

in even dimensions.

From (2.62), the Bogoliubov coefficient $\alpha(t_1;t_0)$ can be found to have the asymptotic form

---

[14] The asymptotic forms given in Eq. (4.67) do not satisfy the normalization condition (2.61). We actually need terms proportional to $\big(1-t_1^2\big)^{\nu_\varepsilon/2}$, which are omitted in deriving the asymptotic forms, in order to ensure the condition. However, these asymptotic forms are yet sufficient for calculating various propagators.

$$\begin{aligned}\alpha(t_1;t_0)\sim\; & -\frac{2\mathrm{i}\sin(\pi\nu_\varepsilon)}{\pi}\left[\frac{2^{\nu_\varepsilon-1}\,\Gamma(\nu_\varepsilon)}{\sqrt{2\bar{m}_1}}\Big(-\frac{d-1}{2}+\nu_\varepsilon+\mathrm{i}\bar{m}_1\Big)\,\big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}}\right]\\ & \times\left[-\frac{2^{\nu_\varepsilon-1}\,\cos(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\sqrt{2\bar{m}_0}\Gamma(k+\nu_\varepsilon+1)}\,\Big(\frac{d-1}{2}-\nu_\varepsilon-\mathrm{i}\bar{m}_0\Big)\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}}\right], \quad (4.72)\\ \beta(t_1;t_0)\sim\; & -\frac{2\mathrm{i}\sin(\pi\nu_\varepsilon)}{\pi}\left[\frac{2^{\nu_\varepsilon-1}\,\Gamma(\nu_\varepsilon)}{\sqrt{2\bar{m}_1}}\Big(-\frac{d-1}{2}+\nu_\varepsilon-\mathrm{i}\bar{m}_1\Big)\big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}}\right]\\ & \times\left[-\frac{2^{\nu_\varepsilon-1}\cos(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\sqrt{2\bar{m}_0}\,\Gamma(k+\nu_\varepsilon+1)}\Big(\frac{d-1}{2}-\nu_\varepsilon+\mathrm{i}\bar{m}_0\Big)\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}}\right] \quad (4.73)\end{aligned}$$

in odd dimensions, and

$$\begin{aligned}\alpha(t_1;t_0)\sim\; & -\mathrm{i}\cos(\pi\nu_\varepsilon)\left[\frac{2^{\nu_\varepsilon-1}\,\Gamma(\nu_\varepsilon)}{\sqrt{2\bar{m}_1}}\Big(-\frac{d-1}{2}+\nu_\varepsilon+\mathrm{i}\bar{m}_1\Big)\,\big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}}\right]\\ & \times\left[\frac{2^{\nu_\varepsilon}\,\sin(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\pi\,\sqrt{2\bar{m}_0}\,\Gamma(k+\nu_\varepsilon+1)}\,\Big(\frac{d-1}{2}-\nu_\varepsilon-\mathrm{i}\bar{m}_0\Big)\,\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}}\right], \quad (4.74)\\ \beta(t_1;t_0)\sim\; & -\frac{2\mathrm{i}\sin(\pi\nu_\varepsilon)}{\pi}\left[\frac{2^{\nu_\varepsilon-1}\,\Gamma(\nu_\varepsilon)}{\sqrt{2\bar{m}_1}}\Big(-\frac{d-1}{2}+\nu_\varepsilon-\mathrm{i}\bar{m}_1\Big)\big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}}\right]\\ & \times\left[\frac{2^{\nu_\varepsilon}\sin(\pi k)\,\Gamma(\nu_\varepsilon)\,\Gamma(k-\nu_\varepsilon+1)}{\pi\sqrt{2\bar{m}_0}\,\Gamma(k+\nu_\varepsilon+1)}\Big(\frac{d-1}{2}-\nu_\varepsilon+\mathrm{i}\bar{m}_0\Big)\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}}\right] \quad (4.75)\end{aligned}$$

in even dimensions.

Although $\varphi(t;t_1)$ and $\bar{\varphi}(t;t_0)$ asymptotically diverge as $\big(1-t_1^2\big)^{-\frac{\nu_\varepsilon}{2}}$ and $\big(1-t_0^2\big)^{-\frac{\nu_\varepsilon}{2}}$, respectively, in the limit $t_1\to+1$ and $t_0\to-1$, we still obtain a finite result for the in-out propagator since the divergence cancels out with that of $\alpha(t_1;t_0)$ (as in the case of the Poincaré patch):

$$\begin{aligned}G_L^{\mathrm{in/out}}(t,t') &= \lim_{\substack{t_0\to-1\\ t_1\to+1}}\frac{1}{\alpha(t_1;t_0)}\,\varphi(t_>;t_1)\,\bar{\varphi}(t_<;t_0)\\ &= \begin{cases}\dfrac{\mathrm{i}\pi}{2\sin\pi\nu_\varepsilon}\,\big[(1-t_>^2)\,(1-t_<^2)\big]^{(d-1)/4}\,\mathsf{P}_k^{-\nu_\varepsilon}(t_>)\,\mathsf{P}_k^{\nu_\varepsilon}(t_<) & (d:\text{odd})\,,\\ \dfrac{\mathrm{i}}{\cos\pi\nu_\varepsilon}\,\big[(1-t_>^2)\,(1-t_<^2)\big]^{(d-1)/4}\,\mathsf{P}_k^{-\nu_\varepsilon}(t_>)\,\mathsf{Q}_k^{\nu_\varepsilon}(t_<) & (d:\text{even})\,.\end{cases}\end{aligned} \quad (4.76)$$

Once this expression is obtained, we may take the limit $\varepsilon\to 0$.

On the other hand, the in-in propagator takes the form

$$\begin{aligned}& G_L^{\mathrm{in/in}}(t,t')\\ &= \lim_{t_0\to-1}\frac{\mathrm{i}}{W_\rho[\bar{\varphi}^*(t;t_0),\,\bar{\varphi}(t;t_0)](T_s)}\,\bar{\varphi}^*(t_>;t_0)\,\bar{\varphi}(t_<;t_0)\end{aligned}$$

$$
= \begin{cases} \dfrac{\mathrm{i}(1-T_s^2)^{-1}}{W_\rho[\mathsf{P}_k^{\nu_\varepsilon^*}(t), \mathsf{P}_k^{\nu_\varepsilon}(t)](T_s)} \left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \mathsf{P}_k^{\nu_\varepsilon^*}(t_>)\, \mathsf{P}_k^{\nu_\varepsilon}(t_<) \quad (d:\mathrm{odd})\,, \\ \dfrac{\mathrm{i}(1-T_s^2)^{-1}}{W_\rho[\mathsf{Q}_k^{\nu_\varepsilon^*}(t), \mathsf{Q}_k^{\nu_\varepsilon}(t)](T_s)} \left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \mathsf{Q}_k^{\nu_\varepsilon^*}(t_>)\, \mathsf{Q}_k^{\nu_\varepsilon}(t_<) \quad (d:\mathrm{even})\,. \end{cases} \tag{4.77}
$$

When the mass is large ($m \geq (d-1)/2$; $\nu_\varepsilon = \mathrm{i}\mu + \varepsilon$), the in-in propagator has a finite limit as $\varepsilon \to 0$:

$$
G_L^{\mathrm{in/in}}(t,t') = \begin{cases} \dfrac{\pi}{2\sinh(\pi\mu)} \left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \mathsf{P}_k^{-\mathrm{i}\mu}(t_>)\, \mathsf{P}_k^{\mathrm{i}\mu}(t_<) \quad (d:\mathrm{odd})\,, \\ \dfrac{2}{\pi\sinh(\pi\mu)} \left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \mathsf{Q}_k^{-\mathrm{i}\mu}(t_>)\, \mathsf{Q}_k^{\mathrm{i}\mu}(t_<) \quad (d:\mathrm{even})\,. \end{cases} \tag{4.78}
$$

In contrast, when the mass is small ($m < (d-1)/2$; $\nu_\varepsilon = \nu + \mathrm{i}\varepsilon$), the Wronskian in the denominator of (4.77) vanishes as $\varepsilon \to 0$, and the in-in propagator diverges.

The wave functions at the remote past and future had been obtained for the heavy mass case ($m \geq (d-1)/2$) in [2, 5] as[15]

$$
\varphi_{\mathrm{out}}(t) \propto (1-t^2)^{\frac{d-1}{4}}\, \mathsf{P}_k^{-\mathrm{i}\mu}(t)\,, \tag{4.79}
$$

$$
\varphi_{\mathrm{in}}^*(t) \propto \begin{cases} (1-t^2)^{\frac{d-1}{4}}\, \mathsf{P}_k^{\mathrm{i}\mu}(t) \quad (d:\mathrm{odd}) \\ (1-t^2)^{\frac{d-1}{4}}\, \mathsf{Q}_k^{\mathrm{i}\mu}(t) \quad (d:\mathrm{even}) \end{cases}, \tag{4.80}
$$

by requiring that $\varphi_{\mathrm{in}}^*(t)$ ($\varphi_{\mathrm{out}}(t)$) is regular for $t \to -1$ ($t \to +1$) and an analytic function in the lower half of complex m^2 plane (see also [13] where they are obtained by suitably choosing the Jost functions). Our propagators (4.76) and (4.77) for $m \geq (d-1)/2$ are consistent with these wave functions.

4.3.2. Propagators in the global patch

We now take the summation over all modes to obtain the de Sitter invariant propagators, $G^{\mathrm{in/out}}(x,x')$ and $G^{\mathrm{in/in}}(x,x')$. For $d \geq 3$, the summation over M can be written with the

---

[15] We here give the following identities which are useful in comparing our results with the literature:

$$
\frac{2^{\frac{d-1}{2}+L}\,\cosh^L\tau\, e^{(-\frac{d-1}{2}-L-\nu)\,\tau}}{\Gamma(1+\nu)}\, F\Big(\frac{d-1}{2}+L, \frac{d-1}{2}+L+\nu; 1+\nu; -e^{-2\tau}\Big) = \big(1-t^2\big)^{\frac{d-1}{4}}\, \mathsf{P}_k^{-\nu}(t)\,,
$$

$$
\begin{aligned} \frac{2^{\frac{d-1}{2}+L}\,\cosh^L\tau\, e^{(\frac{d-1}{2}+L-\nu)\,\tau}}{\Gamma(1-\nu)}\, F\Big(\frac{d-1}{2}+L, \frac{d-1}{2}+L-\nu; 1-\nu; -e^{+2\tau}\Big) &= \big(1-t^2\big)^{\frac{d-1}{4}}\, \mathsf{P}_k^{\nu}(-t) \\ &= \frac{\Gamma(k+1+\nu)}{\Gamma(k+1-\nu)}\,(1-t^2)^{\frac{d-1}{4}} \begin{cases} (-1)^k\, \mathsf{P}_k^{-\nu}(t) & (d:\mathrm{odd}) \\ (-1)^{k+1/2}\,(2/\pi)\, \mathsf{Q}_k^{-\nu}(t) & (d:\mathrm{even}) \end{cases}. \end{aligned}
$$

Gegenbauer polynomials:

$$\sum_{M=1}^{N_L^{(d)}} Y_{LM}(\mathbf{\Omega})\, Y_{LM}(\mathbf{\Omega}') = \frac{2L+d-2}{(d-2)\,|\Omega_{d-1}|}\, C_L^{\frac{d-2}{2}}(\mathbf{\Omega}\cdot\mathbf{\Omega}') \quad \bigl(|\Omega_{d-1}| = 2\pi^{\frac{d}{2}}/\Gamma(d/2)\bigr)\,. \tag{4.81}$$

As for $d = 2$, the sum has the form

$$\sum_{M=1}^{N_L^{(2)}} Y_{LM}(\mathbf{\Omega})\, Y_{LM}(\mathbf{\Omega}') = \begin{cases} \dfrac{1}{2\pi} & (L=0)\,, \\ \dfrac{\cos(L\mathbf{\Omega}\cdot\mathbf{\Omega})}{\pi} & (L\geq 1)\,, \end{cases} \tag{4.82}$$

which is the same as the $d \to 2$ limit of the expression (4.81). Thus Eq. (4.81) can be understood to hold for any dimensionality $d \geq 2$. The in-out propagator in spacetime then takes the form

$$\begin{aligned} G^{\text{in/out}}(x,x') &= \sum_{L=0}^{\infty} G_L^{\text{in/out}}(t,t') \sum_{M=1}^{N_L^{(d)}} Y_{LM}(\mathbf{\Omega})\, Y_{LM}(\mathbf{\Omega}') \\ &= \sum_{L=0}^{\infty} \frac{2L+d-2}{(d-2)\,|\Omega_{d-1}|}\, G_L^{\text{in/out}}(t,t')\, C_L^{\frac{d-2}{2}}(\mathbf{\Omega}\cdot\mathbf{\Omega}')\,, \end{aligned} \tag{4.83}$$

which becomes

$$\begin{aligned} G_{\text{odd}}^{\text{in/out}}(x,x') = &\frac{\mathrm{i}\pi}{2\sin(\pi\nu)\,(d-2)\,|\Omega_{d-1}|} \left[(1-t_>^2)\,(1-t_<^2)\right]^{(d-1)/4} \\ &\times \sum_{L=0}^{\infty} (2L+d-2)\, \mathsf{P}_k^{-\nu}(t_>)\, \mathsf{P}_k^{\nu}(t_<)\, C_L^{\frac{d-2}{2}}(\cos\theta) \end{aligned} \tag{4.84}$$

in odd dimensions, and

$$\begin{aligned} G_{\text{even}}^{\text{in/out}}(x,x') = &\frac{\mathrm{i}}{\cos(\pi\nu)\,(d-2)\,|\Omega_{d-1}|} \left[(1-t_>^2)\,(1-t_<^2)\right]^{(d-1)/4} \\ &\times \sum_{L=0}^{\infty} (2L+d-2)\, \mathsf{P}_k^{-\nu}(t_>)\, \mathsf{Q}_k^{\nu}(t_<)\, C_L^{\frac{d-2}{2}}(\cos\theta) \end{aligned} \tag{4.85}$$

in even dimensions. Here, we have defined θ via the relation $\mathbf{\Omega}\cdot\mathbf{\Omega}' \equiv \cos\theta$.

Using Eqs. (E.25) and (E.26) and introducing

$$u_\pm(x,x') \equiv -Z(x,x') \pm \mathrm{i}0 = -\frac{-t\,t' + \cos\theta}{(1-t^2)^{\frac{1}{2}}\,(1-t'^2)^{\frac{1}{2}}} \pm \mathrm{i}0\,, \tag{4.86}$$

we can rewrite the in-out propagator in a de Sitter invariant form:

$$G^{\mathrm{in/out}}_{\mathrm{odd}}(x,x') = \frac{-e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\sin(\pi\nu)}\left[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) - (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-)\right], \tag{4.87}$$

$$G^{\mathrm{in/out}}_{\mathrm{even}}(x,x') = \frac{\mathrm{i}e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\cos(\pi\nu)}\left[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) + (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-)\right]. \tag{4.88}$$

In the massless limit $m \to 0$ (or $\nu \to (d-1)/2$), we have

$$\sin\pi\nu \to 0 \quad (d:\mathrm{odd})\,, \qquad \cos\pi\nu \to 0 \quad (d:\mathrm{even})\,, \tag{4.89}$$

and thus the propagators (4.87) and (4.88) diverge. We thus conclude that there exists no finite massless limit in the global patch, as opposed to the case of the Poincaré patch.

On the other hand, the in-in propagator in the heavy mass case ($m > (d-1)/2$) takes the form

$$G^{\mathrm{in/in}}(x,x') = \sum_{L=0}^{\infty} \frac{2L+d-2}{(d-2)\,|\Omega_{d-1}|}\,G^{\mathrm{in/in}}_L(t,t')\,C_L^{\frac{d-2}{2}}(\boldsymbol{\Omega}\cdot\boldsymbol{\Omega}')\,, \tag{4.90}$$

which becomes

$$\begin{aligned} G^{\mathrm{in/in}}_{\mathrm{odd}}(x,x') = {}& \frac{\mathrm{i}\pi}{2\sin(\pi\nu)\,(d-2)\,|\Omega_{d-1}|}\left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \\ & \times \sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(t_>)\,\mathsf{P}_k^{\nu}(t_<)\,C_L^{\frac{d-2}{2}}(\cos\theta) \end{aligned} \tag{4.91}$$

in odd dimensions, and

$$\begin{aligned} G^{\mathrm{in/in}}_{\mathrm{even}}(x,x') = {}& \frac{2\mathrm{i}}{\pi\sin(\pi\nu)\,(d-2)\,|\Omega_{d-1}|}\left[(1-t_>^2)\,(1-t_<^2)\right]^{\frac{d-1}{4}} \\ & \times \sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{Q}_k^{-\nu}(t_>)\,\mathsf{Q}_k^{\nu}(t_<)\,C_L^{\frac{d-2}{2}}(\cos\theta) \end{aligned} \tag{4.92}$$

in even dimensions. Here, $\nu = \mathrm{i}\mu = \mathrm{i}\sqrt{m^2-(d-1)^2/4}$ ($\mu \in \mathbb{R}_+$). Note that $G^{\mathrm{in/in}}_{\mathrm{odd}}(x,x') = G^{\mathrm{in/out}}_{\mathrm{odd}}(x,x')$, which is consistent with a well-known fact that the in-vacuum equals the out-vacuum up to a phase in odd dimensions (see [5]). The summations in (4.91) and (4.92) can be carried out analytically by using (E.25) and (E.28), and are again expressed in the de

Sitter invariant forms,

$$
\begin{aligned}
&G^{\rm in/in}_{\rm odd}(x,x') = G^{\rm in/out}_{\rm odd}(x,x') \\
&= \frac{-\,e^{-{\rm i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\sin(\pi\nu)}\Big[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) - (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-)\Big]\,, \qquad (4.93)
\end{aligned}
$$

$$
\begin{aligned}
&G^{\rm in/in}_{\rm even}(x,x') \\
&= \frac{e^{-{\rm i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\sin\pi\nu}\Big\{(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) - (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-) \\
&\qquad + \frac{{\rm i}\,\pi}{\cos\pi\nu}\big[e^{-{\rm i}\pi\nu}\,(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) + e^{{\rm i}\pi\nu}\,(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-)\big]\Big\} \\
&= -\,\frac{e^{-{\rm i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\sin\pi\nu}\Big\{(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_+) - (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(u_-) \\
&\qquad - \frac{\pi}{\cos(\pi\nu)}\big[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-u_+) - (u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-u_-)\big]\Big\}\,. \qquad (4.94)
\end{aligned}
$$

5. Feynman path integral in de Sitter space

In this section, we consider the Feynman propagator obtained by the path-integral,

$$
\begin{aligned}
&\langle\phi(x)\,\phi(x')\rangle \equiv \frac{\int[{\rm d}\phi]\,\phi(x)\,\phi(x')\,e^{{\rm i}S_\varepsilon[\phi]}}{\int[{\rm d}\phi]\,e^{{\rm i}S_\varepsilon[\phi]}} \\
&\text{with}\ \ S_\varepsilon[\phi] \equiv -\,\frac{1}{2}\int {\rm d}^dx\,\sqrt{-g}\,\big(g^{\mu\nu}\,\partial_\mu\phi\,\partial_\nu\phi + (m^2-{\rm i}\varepsilon)\,\phi^2\big)\,. \qquad (5.1)
\end{aligned}
$$

We assume that the base space where $\phi(x)$ lives has a sufficiently large noncompact region in the temporal direction near the future and past boundaries at $t = t_f$ and $t = t_i$, respectively (see Fig. 2). Then, due to the existence of the iε term, when sending the initial time t_0 and the final time t_1 (in the Hamiltonian formalism) to the infinite past at t_i and the



Figure 2: The spacetime region where the path integral is performed.

infinite future at t_f, respectively, the initial and final state would be well kept subject to the projection to the instantaneous ground state at each moment t_0 or t_1 . It is thus natural to expect that the propagator defined in (5.1) coincides with the in-out propagator constructed in the preceding sections. In the following, we give an evidence for this assertion by showing that, for each mode, the propagators numerically calculated from (5.1) coincide with our in-out propagators.[16]

We first expand $\phi(x)$ as in (2.5):

$$\phi(x) = \sum_n \phi_n(t)\, Y_n(\boldsymbol{x})\,. \tag{5.2}$$

The propagator is then written as the sum of the propagators of the mode n :

$$\begin{aligned} \langle \phi(x)\, \phi(x') \rangle &= \sum_n \langle \phi_n(t)\, \phi_n(t') \rangle\, Y_n(\boldsymbol{x})\, Y_n(\boldsymbol{x}')\,, \qquad (5.3)\\ \langle \phi_n(t)\, \phi_n(t') \rangle &\equiv \frac{\int [\mathrm{d}\phi_n]\, e^{\mathrm{i} S_{n,\varepsilon}[\phi_n]}\, \phi_n(t)\, \phi_n(t')}{\int [\mathrm{d}\phi_n]\, e^{\mathrm{i} S_{n,\varepsilon}[\phi_n]}}\,,\\ \text{with}\quad S_{n,\varepsilon}[\phi_n] &\equiv \int_{t_i}^{t_f} \mathrm{d}t\, \frac{1}{2}\, \phi_n(t)\, \bigl[-\partial_t \circ \rho \circ \partial_t - \omega^2_{n,\varepsilon}(t) \bigr]\, \phi_n(t)\,. \qquad (5.4) \end{aligned}$$

We redefine the variable $\phi_n(t)$ by $\chi_n(t) = \rho^{-1/2}(t)\, \phi_n(t) \equiv e^{\sigma(t)}\, \phi_n(t)$ and omit the mode label n for simplicity. Then, the action for each mode has the form

$$S_\varepsilon[\chi] = \int_{t_i}^{t_f} \mathrm{d}t\, \frac{1}{2}\, \chi(t)\, \bigl[-\partial_t^2 - \Omega_\varepsilon^2(t) \bigr]\, \chi(t)\,, \tag{5.5}$$

$$\Omega_\varepsilon^2(t) \equiv \omega_\varepsilon^2(t) - \bigl(\dot\sigma(t) \bigr)^2 - \ddot\sigma(t)\,, \tag{5.6}$$

and the propagator for each mode is given by

$$\langle \phi(t)\, \phi(t') \rangle = \bigl[\rho(t)\, \rho(t') \bigr]^{-1/2} \langle \chi(t)\, \chi(t') \rangle = \bigl[\rho(t)\, \rho(t') \bigr]^{-1/2} \bigl\langle t \bigr| \frac{\mathrm{i}}{-\partial_t^2 - \Omega_\varepsilon^2} \bigl| t' \bigr\rangle\,. \tag{5.7}$$

We numerically evaluate the propagator (5.7) by dividing the interval (t_i, t_f) into N parts and by calculating the inverse of the matrix corresponding to $\mathrm{i}^{-1}\, (-\partial_t^2 - \Omega_\varepsilon^2)$. We take a uniform spacing $a \equiv (t_f - t_i)/N$ for brevity and write the time variable as $t = a\, r$ with r an integer in the region $r_i < r < r_f$ ($r_i \equiv t_i/a$ and $r_f \equiv t_f/a$). We then introduce dimensionless variables χ_r as

$$\chi(t) = a^{1/2}\, \chi_r\,, \tag{5.8}$$

---

[16] Note that one does not need to specify the foliation in defining the path integral (5.1), and the obtained propagator must be de Sitter invariant as long as the symmetry is not broken by any of the UV and IR regulators (which are absent in free field theory considered here). The de Sitter invariance of the in-out propagators obtained in the previous section is an expected property if this equivalence is correct.

with which the action becomes

$$S_\varepsilon[\chi] = \frac{1}{2}\sum_{r,s} S_{\varepsilon,rs}\,\chi_r\,\chi_s\,, \tag{5.9}$$

$$S_{\varepsilon,rs} \equiv \bigl[2 - a^2\,\Omega_\varepsilon(ar)\bigr]\,\delta_{r,s} - \delta_{r,\,s+1} - \delta_{r,\,s-1}\,. \tag{5.10}$$

Since

$$\langle \chi_r \chi_{r'} \rangle = \frac{\int \mathrm{d}^N\chi\ \chi_r\chi_{r'}\exp\bigl(\frac{\mathrm{i}}{2}\sum S_{\varepsilon,\,ss'}\,\chi_s\,\chi_{s'}\bigr)}{\int \mathrm{d}^N\chi\ \ \exp\bigl(\frac{\mathrm{i}}{2}\sum S_{\varepsilon,\,ss'}\,\chi_s\,\chi_{s'}\bigr)} = \mathrm{i}(S_\varepsilon^{-1})_{rr'}\,, \tag{5.11}$$

the propagator is obtained as

$$\langle \phi(t)\,\phi(t')\rangle = \mathrm{i}a\,\bigl[\rho(t)\,\rho(t')\bigr]^{-1/2}\,\bigl(S_\varepsilon^{-1}\bigr)_{r\,r'} \quad \bigl(t = a\,r\,,\ t' = a\,r'\bigr)\,. \tag{5.12}$$

We numerically calculate the inverse matrix (5.12) for both of the Poincaré and global patches and compare the result with our in-out propagators obtained in section 4.

The result in the Poincaré case is depicted in Fig. 3, while that in the global patch of $d = 3$ ($d = 4$) is in Fig. 4 (Fig. 5). We find that there is a perfect agreement for the global patch, while there exists a small discrepancy for the Poincaré patch. We observe that the discrepancy gets reduced as one takes a finer mesh near $\eta = 0$ and a larger value for $|\eta_0|$, and expect that it will disappear eventually. We thus are almost convinced that our in-out propagators coincide with those obtained by the Feynman path integral with the iε prescription.



Figure 3: The real part (left) and the imaginary part (right) of the in-out propagator $G_{\boldsymbol{k}}^{\text{in/out}}(\eta,\eta')$ (solid curve) and the propagator $\langle \phi_{\boldsymbol{k}}(\eta)\,\phi_{\boldsymbol{k}}(\eta')\rangle$ (dashed curve) for $d = 3$, $m = 4$, $k = 1$, $a = 0.002$, $\varepsilon = 0.8$, $\eta_0 = -20$, $\eta_1 = -0.01$, and $\eta' = -10.005$. Recall that $\eta_i = -\infty$ and $\eta_f = 0$.



Figure 4: The real part (left) and the imaginary part (right) of the in-out propagator $G_L^{\mathrm{in/out}}(t,0)$ (solid curve) and the propagator $\langle \phi_L(t)\, \phi_L(0)\rangle$ (dashed curve) for $d=3$, $m=0.5$, $L=1$, $a=0.005$, $\varepsilon=10^{-10}$, $t_0=-0.995$, and $t_1=0.995$. Recall that $t_i=-1$ and $t_f=+1$.



Figure 5: The real part (left) and the imaginary part (right) of the in-out propagator $G_L^{\mathrm{in/out}}(t,0)$ (solid curve) and the propagator $\langle \phi_L(t)\, \phi_L(0)\rangle$ (dashed curve) for $d=4$, $m=0.5$, $L=1$, $a=0.005$, $\varepsilon=10^{-10}$, $t_0=-0.995$, and $t_1=0.995$. Recall that $t_i=-1$ and $t_f=+1$.

6. Heat kernel representation and the composition principle

6.1. General theory

We consider the random walk of a relativistic particle moving in a Lorentzian manifold with the metric

$$\mathrm{d}s^2 = g_{\mu\nu}(x)\,\mathrm{d}x^\mu\,\mathrm{d}x^\nu\,. \tag{6.1}$$

Its trajectory is uniquely specified by the functions $X^\mu(\lambda)$ $(0\leq\lambda\leq 1)$, up to reparametrizations $\lambda\to f(\lambda)$ such that $\mathrm{d}f(\lambda)/\mathrm{d}\lambda > 0,\ f(0)=0,\ f(1)=1$. The amplitude connecting two points x and x' is then given by the Feynman path integral:

$$A(x,x') = \int_{X(0)=x'}^{X(1)=x} \frac{[\mathrm{d}X^\mu(\lambda)]}{\mathrm{Vol(Diff}_1)}\, e^{\,\mathrm{i}I_0[X(\lambda)]}\,, \tag{6.2}$$

where $\mathrm{Vol(Diff}_1)$ is the gauge volume of one-dimensional diffeomorphisms

$$X^\mu(\lambda)\to\tilde{X}^\mu(\lambda) = X^\mu(f(\lambda))\,, \tag{6.3}$$

and we propose to set the action $I_0[X]$ for the random walk in a Lorentzian manifold as

$$I_0\bigl[X(\lambda)\bigr] \equiv -(m-\mathrm{i}\varepsilon)\int_0^1 \mathrm{d}\lambda\,\sqrt{-g_{\mu\nu}\bigl(X(\lambda)\bigr)\,\dot{X}^\mu(\lambda)\,\dot{X}^\nu(\lambda) - \mathrm{i}\varepsilon'}\,. \tag{6.4}$$

Note the presence of two infinitesimal imaginary parts, $\mathrm{i}\varepsilon$ and $\mathrm{i}\varepsilon'$, in $I_0[X(\lambda)]$ $(\varepsilon,\,\varepsilon'>0)$. The first $(\mathrm{i}\varepsilon)$ is the standard one, which manifestly suppresses the contribution from such paths that are prolonged in the timelike direction. We further have introduced the second one $(\mathrm{i}\varepsilon')$ in order to define the path integral for any shape of path in a Lorentzian manifold. In fact, for a timelike segment $(\dot{X}^2<0)$ we can neglect ε' and the action becomes the standard action for a timelike path, while for a spacelike segment $(\dot{X}^2>0)$ we can rewrite the square root as

$$\sqrt{-\dot{X}^2-\mathrm{i}\varepsilon'} = \sqrt{e^{-\mathrm{i}(\pi-0)}\dot{X}^2} = -\,\mathrm{i}\sqrt{\dot{X}^2} \tag{6.5}$$

and the action gives the path-integral weight which suppresses the contribution from such paths that are stretched largely in the spacelike direction.

The action (6.4) of Nambu-Goto type is equivalent to the following action of Polyakov type:

$$I[X(\lambda),e(\lambda)] = \int_0^1 \mathrm{d}\lambda\,\Bigl[\,\frac{\dot{X}^2+\mathrm{i}\varepsilon'}{2\,e} - \frac{m^2-\mathrm{i}\varepsilon}{2}\,e\,\Bigr]\,, \tag{6.6}$$

where $e(\lambda) > 0$ is the einbein defined on the one-dimensional manifold. We can see from this expression that ε and ε' give imaginary parts of the same sign. The new action $I[X(\lambda), e(\lambda)]$ also has the invariance under the one-dimensional diffeomorphisms

$$X^\mu(\lambda) \to \tilde{X}^\mu(\lambda) = X^\mu(f(\lambda))\,, \quad e(\lambda) \to \tilde{e}(\lambda) = \frac{\mathrm{d}f(\lambda)}{\mathrm{d}\lambda}\, e(f(\lambda))\,, \tag{6.7}$$

and we can take a gauge fixing where $e(\lambda) = \text{constant} \equiv T$. However, as is discussed in detail in [14], such constant $T = \int_0^1 \mathrm{d}\lambda\, e(\lambda)$ is actually Diff_1-invariant and needs to be further integrated, so that we obtain[17]

$$\begin{aligned} A(x,x') &= \int_{X(0)=x'}^{X(1)=x} \frac{[\mathrm{d}X^\mu(\lambda)\, \mathrm{d}e(\lambda)]}{\mathrm{Vol}(\mathrm{Diff}_1)}\, e^{\,\mathrm{i} I_0[X(\lambda)]} \\ &= \int_0^\infty \mathrm{d}T \int_{X(0)=x'}^{X(1)=x} [\mathrm{d}X(\lambda)]\, \exp\Big[\,\mathrm{i} \int_0^1 \mathrm{d}\lambda \Big(\frac{\dot{X}^2 + \mathrm{i}\varepsilon'}{2T} - \frac{m^2 - \mathrm{i}\varepsilon}{2}\, T \Big)\Big] \\ &= \int_0^\infty \mathrm{d}T\, e^{-\varepsilon'/(2T)} \int_{X(0)=x'}^{X(T)=x} [\mathrm{d}X(t)]\, \exp\Big[\,\mathrm{i} \int_0^T \mathrm{d}t \Big(\frac{\dot{X}^2(t) + \mathrm{i}\varepsilon'}{2} - \frac{m^2 - \mathrm{i}\varepsilon}{2} \Big)\Big]\,, \end{aligned} \tag{6.8}$$

where in the last line we have rewritten the expression with $t \equiv T\lambda$. The path integral is nothing but that for the quantum mechanical amplitude from the state $|x'\rangle$ to the state $|x\rangle$ with the Hamiltonian

$$H = \frac{1}{2}\big(-\Box + m^2 - \mathrm{i}\varepsilon\big) \equiv \frac{1}{2}\Big[-\frac{1}{\sqrt{-g}}\,\partial_\mu\left(\sqrt{-g}\, g^{\mu\nu}\,\partial_\nu\right) + m^2 - \mathrm{i}\varepsilon\Big]\,, \tag{6.9}$$

and thus we obtain the expression

$$\begin{aligned} A(x,x') &= \int_0^\infty \mathrm{d}T\, e^{-\varepsilon'/(2T)}\, \langle x| e^{-\mathrm{i}(T/2)\,(-\Box + m^2 - \mathrm{i}\varepsilon)} |x'\rangle \\ &= \int_0^\infty \mathrm{d}T\, e^{-\mathrm{i}(T/2)\,(m^2 - \mathrm{i}\varepsilon) - \varepsilon'/(2T)}\, K(x,x';T)\,. \end{aligned} \tag{6.10}$$

Here, $K(x,x';T)$ is the heat kernel of the d'Alembertian $\Box$,

$$K(x,x';T) \equiv \langle x| e^{\mathrm{i}(T/2)\,\Box} |x'\rangle\,, \tag{6.11}$$

which satisfies the following equations:

$$\mathrm{i}\,\frac{\partial}{\partial T}\, K(x,x';T) = -\frac{1}{2}\,\Box_x\, K(x,x';T)\,, \tag{6.12}$$

$$K(x,x';T=0) = \delta^d(x,x') \equiv \frac{1}{\sqrt{-g}}\,\delta^d(x-x')\,. \tag{6.13}$$

---

[17] There may arise a divergence when calculating the Jacobian to obtain the second line, but such divergence should be ultra-local in quantum mechanics (i.e., one-dimensional field theory with a coordinate λ) and can be simply dealt with by an additive renormalization of mass m, as in the Euclidean space considered in [14].

Note that we need to multiply (6.8) by 1/2 to obtain the propagator $G(x,x')$ of a neutral particle (i.e., particle = anti-particle):

$$G(x,x') = \frac{1}{2}\,A(x,x') = \frac{1}{2}\int_0^\infty \mathrm{d}T\, e^{-\mathrm{i}(T/2)\,(m^2-\mathrm{i}\varepsilon)-\varepsilon'/(2T)}\,K(x,x';\,T)\,. \tag{6.14}$$

Since the propagator $G(x,x')$ can formally be written as $G(x,x') = \mathrm{i}\,\langle x|\,(\Box - m^2 + \mathrm{i}\varepsilon)^{-1}\,|x'\rangle$, one can easily show that $G(x,x')$ satisfies the following *composition law* [9]:

$$\frac{\partial}{\partial m^2}\,G(x,x') = \mathrm{i}\,\int\!\sqrt{-g(y)}\,\mathrm{d}^d y\,G(x,y)\,G(y,x')\,, \tag{6.15}$$

which is consistent with the asymptotic form of $G(x,x')$ for large timelike separation with large mass:

$$G(x,x') \sim e^{-\mathrm{i}m\,L(x,x')}\,, \tag{6.16}$$

where $L(x,x')$ is the timelike geodesic distance between x and x' .

As a simple example, we consider a neutral particle propagating in a d-dimensional Minkowski space with the metric

$$\mathrm{d}s^2 = -\,\mathrm{d}t^2 + \mathrm{d}\boldsymbol{x}^2\,. \tag{6.17}$$

Then the heat kernel can be calculated with the momentum representation as

$$\begin{aligned} K(x,x';\,T) &= \int \frac{\mathrm{d}^d p}{(2\pi)^d}\, e^{\mathrm{i}p(x-x')}\, e^{-\mathrm{i}(T/2)\,p^2} \\ &= \mathrm{i}\,\Bigl(\frac{1}{2\pi \mathrm{i}T}\Bigr)^{d/2}\,\exp\Bigl[\mathrm{i}\frac{(x-x')^2}{2T}\Bigr]\,. \end{aligned} \tag{6.18}$$

Here, the first i reflects the fact that the Gaussian integral over p_0 has the opposite sign of quadratic term to that for the other variables p_i ($i = 1,\ldots,d-1$). Substituting this to (6.14), we obtain the following expression:

$$G(x,x') = \frac{\mathrm{i}^{-(d-2)/2}}{(4\pi)^{d/2}}\int_0^\infty \mathrm{d}s\, s^{-\frac{d}{2}}\, e^{-\frac{z^2}{4\,s}-a\,s} \tag{6.19}$$

with $z^2 = \varepsilon' - \mathrm{i}(x-x')^2$ and $a = \mathrm{i}(m^2 - \mathrm{i}\varepsilon)$. This integration can be easily performed, and we obtain

$$\begin{aligned} G(x,x') &= \frac{\mathrm{i}^{-(d-2)/2}}{(4\pi)^{d/2}}\, a^{\frac{d-2}{2}}\, 2^{\frac{d}{2}}\, \bigl(\sqrt{a}\,z\bigr)^{-\frac{d-2}{2}}\, K_{\frac{d-2}{2}}\bigl(\sqrt{a}\,z\bigr) \\ &= \frac{m^{d-2}}{(2\pi)^{d/2}}\, \bigl(m\sqrt{\sigma + \mathrm{i}\varepsilon'}\bigr)^{-\frac{d-2}{2}}\, K_{\frac{d-2}{2}}\bigl(m\sqrt{\sigma + \mathrm{i}\varepsilon'}\bigr)\,, \end{aligned} \tag{6.20}$$

where $\sigma \equiv (x-x')^2$. This certainly agrees with the propagator (3.29) of a real scalar field.

6.2. de Sitter case

In this subsection, we check that the in-out propagators (4.51), (4.87), and (4.88) indeed satisfies the composition principle by giving their heat kernel representations.[18]

Poincaré patch

We start from the following integral representation of the associated Legendre functions (a proof is given in Appendix F):

$$\mathcal{Q}^{\frac{d-2}{2}}_{\nu-1/2}(u) = e^{\mathrm{i}\pi\frac{d-2}{2}} \int_0^\infty \mathrm{d}\lambda \, \frac{\Gamma\bigl(\frac{d-1}{2}+\mathrm{i}\lambda\bigr)\,\Gamma\bigl(\frac{d-1}{2}-\mathrm{i}\lambda\bigr)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)} \, \frac{\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u)}{\nu^2+\lambda^2} \quad \bigl[d\in\mathbb{Z}\,, \quad \mathrm{Re}\,\nu>0\bigr]\,. \tag{6.21}$$

Since ν_ε^2 has a positive imaginary part, we have

$$\begin{aligned} & e^{-\mathrm{i}\pi\frac{d-2}{2}} \bigl(u^2-1\bigr)^{-\frac{d-2}{4}} \, \mathcal{Q}^{\frac{d-2}{2}}_{\nu_\varepsilon-1/2}(u) \\ &= \frac{1}{2\mathrm{i}} \int_0^\infty \mathrm{d}T \int_0^\infty \mathrm{d}\lambda \, \frac{\Gamma\bigl(\frac{d-1}{2}+\mathrm{i}\lambda\bigr)\,\Gamma\bigl(\frac{d-1}{2}-\mathrm{i}\lambda\bigr)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)} \bigl(u^2-1\bigr)^{-\frac{d-2}{4}} \, \mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u) \, e^{\mathrm{i}\frac{T}{2}(\nu_\varepsilon^2+\lambda^2)} \\ &= \frac{\mathrm{i}}{2^{2-\frac{d}{2}}} \int_0^\infty \mathrm{d}T \int_0^\infty \mathrm{d}\lambda \, \frac{\lambda\,\Gamma\bigl(\frac{d-1}{2}\bigr)\sinh(\pi\lambda)}{\sqrt{\pi}\,\cos\bigl[\pi\bigl(\frac{d}{2}-\mathrm{i}\lambda\bigr)\bigr]} \, C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u) \, e^{\mathrm{i}\frac{T}{2}(\nu_\varepsilon^2+\lambda^2)}\,. \end{aligned} \tag{6.22}$$

Here, $u = Z - \mathrm{i}0$, and to obtain the second line we have used the identity

$$\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-\frac{1}{2}}(u) = 2^{-\frac{d-2}{2}} \bigl(u^2-1\bigr)^{\frac{d-2}{4}} \, \frac{\Gamma(d-1)\,\Gamma\bigl(\mathrm{i}\lambda-\frac{d-3}{2}\bigr)}{\Gamma(d/2)\,\Gamma\bigl(\mathrm{i}\lambda+\frac{d-1}{2}\bigr)} \, C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)\,. \tag{6.23}$$

We thus find that the in-out propagator (4.51) in the Poincaré patch has the heat kernel representation of the form

$$G(x,x') = \frac{1}{2}\int_0^\infty \mathrm{d}T \, e^{-\mathrm{i}\frac{m^2-\mathrm{i}\varepsilon}{2}T} \, K(x,x';T)\,, \tag{6.24}$$

$$\begin{aligned} K(x,x';T) &= -\frac{e^{-\mathrm{i}\pi\frac{d-3}{2}}\bigl(u^2-1\bigr)^{-\frac{d-2}{4}}}{(2\pi)^{d/2}} \\ &\qquad \times \int_0^\infty \mathrm{d}\lambda \, \frac{\Gamma\bigl(\frac{d-1}{2}+\mathrm{i}\lambda\bigr)\,\Gamma\bigl(\frac{d-1}{2}-\mathrm{i}\lambda\bigr)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)} \, \mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u) \, e^{\mathrm{i}\frac{T}{2}\left(\lambda^2+(\frac{d-1}{2})^2\right)} \\ &= \frac{e^{-\mathrm{i}\pi\frac{d-3}{2}}\,\Gamma\bigl(\frac{d-1}{2}\bigr)}{2\pi^{\frac{d+1}{2}}} \int_0^\infty \mathrm{d}\lambda \, \frac{\lambda\sinh(\pi\lambda)}{\cos\bigl[\pi\bigl(\frac{d}{2}-\mathrm{i}\lambda\bigr)\bigr]} \, C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u) \, e^{\mathrm{i}\frac{T}{2}\left(\lambda^2+(\frac{d-1}{2})^2\right)}\,. \end{aligned} \tag{6.25}$$

---

[18] See [15] (also [9]) for the direct evaluation of the random walk in de Sitter space, which is based on the heat kernel for Euclidean AdS space obtained in [16].

The above heat kernel certainly satisfies Eqs. (6.12) and (6.13). In fact, Eq. (6.12) can be shown in the following way:

$$\begin{aligned}
&\mathrm{i}\,\frac{\partial}{\partial T}\,K(x,x';\,T)\\
&= -\frac{e^{-\mathrm{i}\pi\frac{d-3}{2}}\,\Gamma\big(\frac{d-1}{2}\big)}{4\pi^{\frac{d+1}{2}}}\int_0^\infty \mathrm{d}\lambda\,\frac{\lambda\,\sinh(\pi\lambda)\,\big(\lambda^2+\big(\frac{d-1}{2}\big)^2\big)}{\cos\big[\pi\big(\frac{d}{2}-\mathrm{i}\lambda\big)\big]}\,C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)\,e^{\mathrm{i}\frac{T}{2}\,\left(\lambda^2+(\frac{d-1}{2})^2\right)}\\
&= -\frac{1}{2}\left[-\big(Z^2-1\big)\,\partial_Z^2K(x,x';\,T)-d\,Z\,\partial_Z K(x,x';\,T)\right]\\
&= -\frac{1}{2}\,\Box_x K(x,x';\,T)\,, \qquad (6.26)
\end{aligned}$$

where we have used the Gegenbauer differential equation

$$\big(1-u^2\big)\,\partial_u^2 C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)-d\,u\,\partial_u C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)-\left[\lambda^2+\left(\frac{d-1}{2}\right)^2\right]C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)=0\,, \qquad (6.27)$$

and the fact that the Klein-Gordon operator for functions of the de Sitter invariant $f(Z)$ can be written as

$$\Box f(Z)=\big(1-Z^2\big)\,\partial_Z^2 f(Z)-d\,Z\,\partial_Z f(Z)\,. \qquad (6.28)$$

The initial condition (6.13) can be shown to hold by using the heat kernel equation and the equality

$$(\Box_x-m^2+\mathrm{i}\varepsilon)G^{\mathrm{in/out}}(x,x')=\frac{\mathrm{i}}{\sqrt{-g}}\,\delta^d(x-x') \qquad (6.29)$$

as follows:

$$\begin{aligned}
\frac{\mathrm{i}}{\sqrt{-g}}\,\delta^d(x-x') &= \frac{1}{2}\int_0^\infty \mathrm{d}T\,e^{-\mathrm{i}\frac{m^2-\mathrm{i}\varepsilon}{2}T}\,(\Box_x-m^2+\mathrm{i}\varepsilon)K(x,x';\,T)\\
&= -\mathrm{i}\int_0^\infty \mathrm{d}T\,\frac{\partial}{\partial T}\left[e^{-\mathrm{i}\frac{m^2-\mathrm{i}\varepsilon}{2}T}\,K(x,x';\,T)\right]\\
&= \mathrm{i}K(x,x';\,0)\,. \qquad (6.30)
\end{aligned}$$

global patch

In a similar way, using Eq. (6.22), we can show that the in-out propagators (4.87) and (4.88) has the heat kernel representation of the form

$$G^{\mathrm{in/out}}_{\left\{\substack{\mathrm{odd}\\ \mathrm{even}}\right\}}(x,x')=\frac{1}{2}\int_0^\infty \mathrm{d}T\,e^{-\mathrm{i}\frac{m^2-\mathrm{i}\varepsilon}{2}T}\,K^{\mathrm{in/out}}_{\left\{\substack{\mathrm{odd}\\ \mathrm{even}}\right\}}(x,x';\,T) \qquad (6.31)$$

with

$$
\begin{aligned}
&K_{\rm odd}^{\rm in/out}(x,x';\,T) \\
&= \frac{(-1)^{\frac{d+1}{2}}\,\Gamma\bigl(\frac{d-1}{2}\bigr)}{4\pi^{\frac{d+1}{2}}\,\sin(\pi\nu)} \int_0^\infty {\rm d}\lambda\,\lambda\,\bigl[C^{\frac{d-1}{2}}_{{\rm i}\lambda-\frac{d-1}{2}}(u_+) - C^{\frac{d-1}{2}}_{{\rm i}\lambda-\frac{d-1}{2}}(u_-)\bigr]\, e^{{\rm i}\frac{T}{2}\left(\lambda^2+(\frac{d-1}{2})^2\right)}\,, \qquad (6.32)\\
&K_{\rm even}^{\rm in/out}(x,x';\,T) \\
&= \frac{(-1)^{\frac{d+2}{2}}\,\Gamma\bigl(\frac{d-1}{2}\bigr)}{4\pi^{\frac{d+1}{2}}\,\cos(\pi\nu)} \int_0^\infty {\rm d}\lambda\,\lambda\,\tanh(\pi\lambda)\,\bigl[C^{\frac{d-1}{2}}_{{\rm i}\lambda-\frac{d-1}{2}}(u_+) + C^{\frac{d-1}{2}}_{{\rm i}\lambda-\frac{d-1}{2}}(u_-)\bigr]\, e^{{\rm i}\frac{T}{2}\left(\lambda^2+(\frac{d-1}{2})^2\right)}\,. \\
&\hfill (6.33)
\end{aligned}
$$

Equations (6.12) and (6.13) also hold for these heat kernels, as can be shown in the same way as above.

6.3. Relation to the Green function in Euclidean AdS space

As has been pointed out in [15], the in-out propagator in the Poincaré patch is directly related to the Green function in Euclidean AdS space through an analytic continuation. In this subsection, we demonstrate this equivalence with a precise numerical constant.

d-dimensional Euclidean AdS space (EAdS$_d$) is defined as the hypersurface in a $(d+1)$-dimensional Minkowski space with the relation

$$
\eta_{MN}\,Y^M\,Y^N = -\,\ell'^2 \quad (M,N=0,\ldots,d)\,, \qquad (6.34)
$$

where ℓ' is called the AdS radius. EAdS$_d$ has two connected components. A frequently used coordinate system which covers only a single connected component is the Poincaré coordinates (z,y^i) $(i=1,\ldots,d-1)$ that are defined by the following embedding:

$$
Y^0 = \frac{\ell'^2+z^2+|\boldsymbol{y}|^2}{2z}\,,\quad Y^i = \ell'\,\frac{y^i}{z}\,,\quad Y^d = \frac{\ell'^2-z^2-|\boldsymbol{y}|^2}{2z}\,. \qquad (6.35)
$$

We here have chosen the component with $z>0$. The metric then takes the form

$$
{\rm d}s^2 = \ell'^2\,\frac{{\rm d}z^2+{\rm d}\boldsymbol{y}\cdot{\rm d}\boldsymbol{y}}{z^2}\,. \qquad (6.36)
$$

The Green function in EAdS space is known to have the following form (see, e.g., [17]):[19]

$$G_{\rm EAdS}(y,y') = \frac{e^{-{\rm i}\pi\,(d-2)/2}}{(2\pi)^{d/2}}\,(Z'^2(y,y')-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{(d-2)/2}_{\nu'-1/2}(Z'(y,y'))\,, \tag{6.37}$$

where $Z'(y,y')$ is the invariant of EAdS space,

$$Z'(y,y') \equiv\,-\ell'^{-2}\,\eta_{MN}\,Y^M(y)\,Y^N(y') = 1 + \frac{(z-z')^2+|\boldsymbol{y}-\boldsymbol{y}'|^2}{2\,z\,z'}\,, \tag{6.38}$$

and

$$\nu' \equiv \sqrt{\Bigl(\frac{d-1}{2}\Bigr)^2 + m^2\,\ell'^2}\,. \tag{6.39}$$

Note that Z' is always larger than unity.

The coordinate system (z,y^i) is related to the Poincaré coordinates (η,x^i) of d-dimensional de Sitter space through the analytic continuation

$$z = e^{{\rm i}\,\frac{\pi-0}{2}}(-\eta)\,,\quad y^i = x^i\,,\quad \ell' = e^{{\rm i}\,\frac{\pi-0}{2}}\,\ell\,, \tag{6.40}$$

or equivalently,

$$Y^0 = {\rm i}X^d\,,\quad Y^i = X^i\,,\quad Y^d = {\rm i}X^0\,. \tag{6.41}$$

In fact, one can easily show that the metrics of EAdS$_d$ and dS$_d$ transform to each other. One also finds the relations

$$\begin{aligned}\nu' &= \sqrt{\Bigl(\frac{d-1}{2}\Bigr)^2 + m^2\,\ell'^2} = \sqrt{\Bigl(\frac{d-1}{2}\Bigr)^2 - m^2\,\ell^2 + {\rm i}0}\\ &= \nu_\varepsilon\,, \end{aligned}\tag{6.42}$$

$$\begin{aligned}Z'(y,y') &= \frac{z^2+z'^2+|\boldsymbol{y}-\boldsymbol{y}'|^2}{2\,z\,z'} = \frac{(-\eta)^2+(-\eta')^2-|\boldsymbol{x}-\boldsymbol{x}'|^2}{2\eta\,\eta'} - {\rm i}0\\ &= Z(x,x') - {\rm i}0\,,\end{aligned}\tag{6.43}$$

with which the Green function on EAdS space can be rewritten as

$$G_{\rm EAdS}(y,y') = \frac{e^{-{\rm i}\pi\,(d-2)/2}}{(2\pi)^{d/2}}\,(u^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{(d-2)/2}_{\nu_\varepsilon-1/2}(u)\quad (u = Z(x,x') - {\rm i}0)\,. \tag{6.44}$$

---

[19] In fact, solving the Klein-Gordon equation with a delta function source using the EAdS invariant $Z' = Z'(y,y')$, we see that the Green function is a linear combination of $(Z'^2-1)^{-(d-2)/4}\mathcal{P}^{(d-2)/2}_{\nu'-1/2}(Z')$ and $(Z'^2-1)^{-(d-2)/4}\mathcal{Q}^{(d-2)/2}_{\nu'-1/2}(Z')$. By requiring that the Green function damps at large separation (cluster property), only the latter solution is selected, as can be seen from the asymptotic forms of the associated Legendre functions [see (E.19) and (E.20)]. The normalization is then determined by requiring that the Green function coincides with that in Euclidean space for infinitesimal separation of y and y' [or $Z'(y,y')\to 1$].

This coincides with the in-out propagator (4.51) in the Poincaré patch of de Sitter space up to a phase:

$$G^{\mathrm{in/out}}(x,x') = \frac{e^{-\mathrm{i}\pi(d-2)}}{(2\pi)^{d/2}}\,(u^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}^{(d-2)/2}_{\nu-1/2}(u) \quad (u = Z(x,x') - \mathrm{i}0)\,. \tag{6.45}$$

As pointed out in [5], the Green function of EAdS space has no direct relation with the in-in propagators associated with α vacuum of de Sitter space for any α. We see that it is the in-out propagator (in the Poincaré patch) which is actually related to the Green function of EAdS space. We thus expect that we can obtain a deep insight on the dS/CFT correspondence [18] by analytically continuing the Euclidean AdS/CFT correspondence and by interpreting the result in terms of the in-out propagators (not of the in-in propagators).

7. Discussions and conclusion

In this paper, we have considered quantum theory of a free scalar field in a nonstatic spacetime. We first developed a framework to treat a harmonic oscillator with time-dependent parameters, and then applied it to investigate a free scalar field in de Sitter space, both in the Poincaré and the global patches.

We have introduced the vacuum state at each moment t_I as the instantaneous ground state of the Hamiltonian at the moment. We developed a calculation method to obtain the wave function $\varphi(t;t_I)$ associated with the vacuum. The in-in and in-out propagators are then obtained from the wave functions by sending the initial and final times to the past and future infinities.

A major advantage of our prescription in defining the vacuum is that we do not need to introduce "positive-energy wave functions" that cannot be defined in a definite way for a spacetime with no asymptotic timelike Killing vector.

We have applied our method to calculate the in-in and in-out propagators in de Sitter space. The obtained propagators take de Sitter invariant forms, and are consistent with the results known in the literature. What actually happens is that, when the time t_I is sent to the temporal boundary, our wave functions $\varphi(t;t_I)$ may have different forms from those of the in- or out-wave functions in the literature, but the obtained propagators yet coincide with each other (see footnote 14 of subsec. 4.3).

As a new result, we have found that a finite massless limit exists for the in-out propagator in the Poincaré patch (with minimal coupling). This is in contrast to the in-in propagator, where the no-go theorem is known that no massless limit exists for the in-in propagators without breaking the de Sitter invariance [3]. The same functional form had been obtained for the in-out propagators without precise numerical coefficients in [9, 15] from other approaches, and the massless case also had been considered in [9]. However, one cannot discuss the existence of a finite massless limit without knowing the precise numerical coefficients. Indeed, our in-out propagator in the global patch diverges in the massless limit just because the numerical coefficient diverges.

We have argued that our in-out propagators should coincide with the Feynman propagators obtained by the path integral with the $\mathrm{i}\varepsilon$ prescription, and have given a numerical evidence for the coincidence. We have also shown that the in-out propagators in both the Poincaré and global patches satisfy Polyakov's composition principle. These observations exhibit the naturalness of the in-out propagators. It should be interesting to study under which conditions the coincidence between the in-out propagators based on the instantaneous vacuum and the propagators derived from the path-integral formalism is realized, and furthermore, to investigate whether the composition principle holds universally for such propagators.

Our in-in propagator in the global patch has a finite value for $m \geq (d-1)/2$, but it diverges for $m < (d-1)/2$. It will be important to compare the in-in propagators with those obtained (numerically) by the path integral of the Schwinger-Keldysh type [11, 12] (see also [19]).

As an important application of our construction, we can consider the Unruh-DeWitt detector working in a finite time interval. In the literature, in order to study the thermal properties of de Sitter space, the Euclidean vacuum has been considered. Since it is a vacuum state at the infinite past in the Poincaré patch, the detectors have to start at the infinite past, which is the situation hard to realize in actual experiments. On the other hand, since our method allows us to construct the vacuum state at a very far but finite past, one may investigate the inherent nonequilibrium property of de Sitter space, by taking account of the corrections to the in-in propagators coming from the effect of such finite-time start (which violates the de Sitter invariance) [20]. It would be also interesting to investigate some physical quantities such as the rate of vacuum decay at finite times on the basis of our formalism.

As another future direction, it should be interesting to apply our method to quantum field theories in spacetimes with horizons, such as spacetimes with black holes and de Sitter space in the static patch. For such a spacetime, one needs to carefully study the consistency of our formalism with boundary conditions at the horizon. It would be also interesting to consider the β-vacua in the rotating Rindler-AdS spacetime [21]. Since the inequivalent spacetime foliations (parameterized by β) cover the same portion of the AdS spacetime, one expects that the propagator is uniquely determined by performing the path integral. However, the in- and out-vacua still has a nontrivial dependence on the parameter β labeling the Hamiltonians. Thus it would be possible to observe which β is preferred by the path-integral formalism.

It should be important to consider interacting fields in generic nonstatic spacetimes and to establish perturbation theory on the basis of our formalism. It will be also interesting to investigate the in-out propagators for gravitons, since our method can be applied to field theory of higher spins without any essential modifications.

Acknowledgments

This work was supported by the Grant-in-Aid for the Global COE program "The Next Generation of Physics, Spun from Universality and Emergence" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. This work was also supported by MEXT (Grant No. 23540304).

A. Proof of Eq. (2.69)

Setting $t = T_s$ in (2.47) and using the hermiticity $q_s^\dagger(T_s) = q_s(T_s)$ and $p_s^\dagger(T_s) = p_s(T_s)$, we obtain

$$\begin{pmatrix} c_1^\dagger \\ c_2^\dagger \end{pmatrix} = \frac{1}{\big(W_\rho[f,g]\big)^*} \begin{pmatrix} W_\rho[f,g^*] & W_\rho[g,g^*] \\ W_\rho[f^*,f] & W_\rho[f^*,g] \end{pmatrix}(T_s) \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \equiv M_s \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} . \tag{A.1}$$

Then, from (2.48) and (2.52), we obtain

$$\begin{pmatrix} a^\dagger(t) \\ \bar{a}^\dagger(t) \end{pmatrix} = [C^{-1}(t)]^* \, M_s \, C(t') \begin{pmatrix} a(t') \\ \bar{a}(t') \end{pmatrix} \equiv \Lambda(t;t') \begin{pmatrix} a(t') \\ \bar{a}(t') \end{pmatrix} . \tag{A.2}$$

A straightforward calculation shows that

$$\Lambda(t;t') = \mathrm{i} \begin{pmatrix} -\,W_\rho[\bar{\varphi}^*(T_s,t),\varphi(T_s,t')] & -\,W_\rho[\bar{\varphi}^*(T_s,t),\bar{\varphi}(T_s,t')] \\ W_\rho[\varphi^*(T_s,t),\varphi(T_s,t')] & W_\rho[\varphi^*(T_s,t),\bar{\varphi}(T_s,t')] \end{pmatrix} . \tag{A.3}$$

Note that $\det\Lambda(t;t') = -1$ due to the commutation relations $[a^\dagger(t), \bar{a}^\dagger(t)] = -1$ and $[a(t'), \bar{a}(t')] = 1$ (this can also be checked by a direct calculation). Then, setting $t = t' = t_0$ in (A.2) and (A.3), we find that

$$\bar{a}_0^\dagger = -\frac{W_\rho[\varphi^*(T_s;t_0), \bar{\varphi}(T_s;t_0)]}{W_\rho[\bar{\varphi}^*(T_s;t_0), \bar{\varphi}(T_s;t_0)]}\, a_0^\dagger + \frac{\mathrm{i}}{W_\rho[\bar{\varphi}^*(T_s;t_0), \bar{\varphi}(T_s;t_0)]}\, a_0\,. \tag{A.4}$$

B. Asymptotically Minkowski space

In this appendix, we investigate a well-studied case where spacetime is asymptotically Minkowski in both the remote past and the remote future.

B.1. Setup

For brevity we consider the two-dimensional spacetime with the metric

$$\mathrm{d}s^2 = a^2(t)\big(-\mathrm{d}t^2 + \mathrm{d}x^2\big)\,, \tag{B.1}$$

where the scale factor $a^2(t)$ now depends on time and takes the form

$$a^2(t) = a_0^2\,\frac{1-\tanh t}{2} + a_1^2\,\frac{1+\tanh t}{2}\,. \tag{B.2}$$

This spacetime is asymptotically Minkowski with scale a_0 in the remote past and with scale a_1 in the remote future. By expanding a scalar field $\phi(t,x)$ as

$$\phi(t,x) = \sum_{k\geq 0}\sum_a \phi_{k,a}(t)\, Y_{k,a}(x) \tag{B.3}$$

as in section 3, the action becomes[20]

$$\begin{aligned} S[\phi(t,x)] &= \int \mathrm{d}t\,\mathrm{d}x\,\sqrt{-g}\,\Big[-\frac{1}{2}\,g^{\mu\nu}\,\partial_\mu\phi\,\partial_\nu\phi - \frac{m^2-\mathrm{i}\varepsilon}{2}\,\phi^2\Big] \\ &= \sum_{k\geq 0}\sum_a \int \mathrm{d}t\,\frac{1}{2}\,\big[\,\dot{\phi}_{k,a}^2(t) - \big(k^2 + m^2\,a^2(t)\big)\,\phi_{k,a}^2(t)\,\big]\,. \end{aligned} \tag{B.4}$$

Thus, the correspondence with the ingredients of section 2 is given by

$$q(t) = \phi_{k,a}(t)\,, \quad \rho(t) = 1\,, \quad \omega(t) = \sqrt{\omega_0^2\,\frac{1-\tanh t}{2} + \omega_1^2\,\frac{1+\tanh t}{2}}\,, \tag{B.5}$$

---

[20] Since $\mathrm{i}\varepsilon$ plays no essential role in this appendix, we have eliminated it from the action.

where

$$\omega_0 \equiv \sqrt{k^2 + m^2\, a_0^2}\,, \quad \omega_1 \equiv \sqrt{k^2 + m^2\, a_1^2}\,. \tag{B.6}$$

We also introduce

$$\omega_\pm \equiv \frac{1}{2}\big(\omega_1 \pm \omega_0\big)\,. \tag{B.7}$$

B.2. Wave functions

The equation of motion takes the form

$$0 = \ddot{q} + \frac{\dot{\rho}}{\rho}\, q + \omega^2\, q = \ddot{q} + \Big(\frac{\omega_1^2 + \omega_0^2}{2} + \frac{\omega_1^2 - \omega_0^2}{2}\, \tanh t\Big)\, q\,, \tag{B.8}$$

which can be solved analytically with the hypergeometric function. We set a pair of independent solutions $\{f(t), g(t)\}$ as

$$f(t) = \Big(\frac{1-\zeta}{2}\Big)^{\mathrm{i}\omega_1/2} \Big(\frac{1+\zeta}{2}\Big)^{-\mathrm{i}\omega_0/2} F\Big(\mathrm{i}\omega_-,\, 1+\mathrm{i}\omega_-\,; 1-\mathrm{i}\omega_0\,; \frac{1+\zeta}{2}\Big)\,, \tag{B.9}$$

$$g(t) = \Big(\frac{1-\zeta}{2}\Big)^{-\mathrm{i}\omega_1/2} \Big(\frac{1+\zeta}{2}\Big)^{\mathrm{i}\omega_0/2} F\Big(-\mathrm{i}\omega_-,\, 1-\mathrm{i}\omega_-\,; 1+\mathrm{i}\omega_0\,; \frac{1+\zeta}{2}\Big)\,, \tag{B.10}$$

where $\zeta \equiv \tanh t$ and $F(a, b; c;\, z)$ is the hypergeometric function. Their asymptotic forms for $t = t_0 \sim -\infty$ (or $\zeta = \zeta_0 \sim -1$) are easily found to be

$$f_0 = f(t_0) \sim e^{-\mathrm{i}\omega_0\, t_0}\,, \quad g_0 = g(t_0) \sim e^{\mathrm{i}\omega_0\, t_0}\,, \tag{B.11}$$

and the weighted Wronskian $W_\rho[f, g] = \rho(t)\, W[f, g](t)$ is found to be

$$W_\rho[f, g] = 2\mathrm{i}\omega_0\,. \tag{B.12}$$

From this we find that the functions in (2.49) have the asymptotic forms

$$u_0 \sim 0\,, \qquad \bar{u}_0 \sim -2\mathrm{i}\omega_0\, e^{-\mathrm{i}\omega_0\, t_0}\,, \tag{B.13}$$

$$v_0 \sim 2\mathrm{i}\omega_0\, e^{\mathrm{i}\omega_0\, t_0}\,, \quad \bar{v}_0 \sim 0\,. \tag{B.14}$$

The wave functions in the limit $t_0 \to -\infty$ then take the form

$$\begin{aligned}\varphi(t; t_0) &\sim \frac{1}{\sqrt{2\omega_0}} e^{\mathrm{i}\omega_0 t_0} f(t) \\ &\sim \frac{1}{\sqrt{2\omega_0}} e^{\mathrm{i}\omega_0 t_0 - \mathrm{i}\omega_+ t - \mathrm{i}\omega_-\, \log(2\cosh t)} F\Big(\mathrm{i}\omega_-,\, 1+\mathrm{i}\omega_-\,; 1-\mathrm{i}\omega_0\,; \frac{1+\tanh t}{2}\Big)\,, \end{aligned} \tag{B.15}$$

$$\begin{aligned}\bar{\varphi}(t; t_0) &\sim \frac{1}{\sqrt{2\omega_0}} e^{-\mathrm{i}\omega_0 t_0} g(t) \\ &\sim \frac{1}{\sqrt{2\omega_0}} e^{-\mathrm{i}\omega_0 t_0 + \mathrm{i}\omega_+ t + \mathrm{i}\omega_-\, \log(2\cosh t)} F\Big(-\mathrm{i}\omega_-,\, 1-\mathrm{i}\omega_-\,; 1+\mathrm{i}\omega_0\,; \frac{1+\tanh t}{2}\Big)\,. \end{aligned} \tag{B.16}$$

In order to calculate the asymptotic forms of various functions for $t \sim +\infty$, it is convenient to rewrite $f(t)$ and $g(t)$ by using the formula

$$\begin{aligned} F(a,b;c;\,z) &= \frac{\Gamma(c)\,\Gamma(c-a-b)}{\Gamma(c-a)\,\Gamma(c-b)}\,F(a,\,b;\,a+b-c+1;\,1-z) \\ &\quad + \frac{\Gamma(c)\,\Gamma(a+b-c)}{\Gamma(a)\,\Gamma(b)}\,(1-z)^{c-a-b}\,F(c-a,\,c-b;\,c-a-b+1;\,1-z)\,. \end{aligned} \tag{B.17}$$

We then obtain

$$\begin{aligned} f(t) &= \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}\Big(\frac{1-\zeta}{2}\Big)^{\mathrm{i}\omega_1/2}\Big(\frac{1+\zeta}{2}\Big)^{-\mathrm{i}\omega_0/2}\,\Big[\tilde{\alpha}^* F\Big(\mathrm{i}\omega_-,\,1+\mathrm{i}\omega_-\,;1+\mathrm{i}\omega_1\,;\frac{1-\zeta}{2}\Big) \\ &\quad - \tilde{\beta}\Big(\frac{1-\zeta}{2}\Big)^{-\mathrm{i}\omega_1} F\Big(1-\mathrm{i}\omega_+,\,-\mathrm{i}\omega_+\,;1-\mathrm{i}\omega_1\,;\frac{1-\zeta}{2}\Big)\Big]\,, \end{aligned} \tag{B.18}$$

$$\begin{aligned} g(t) &= \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}\Big(\frac{1-\zeta}{2}\Big)^{-\mathrm{i}\omega_1/2}\Big(\frac{1+\zeta}{2}\Big)^{\mathrm{i}\omega_0/2}\,\Big[\tilde{\alpha} F\Big(-\mathrm{i}\omega_-,\,1-\mathrm{i}\omega_-\,;1-\mathrm{i}\omega_1\,;\frac{1-\zeta}{2}\Big) \\ &\quad - \tilde{\beta}^*\Big(\frac{1-\zeta}{2}\Big)^{\mathrm{i}\omega_1} F\Big(1+\mathrm{i}\omega_+,\,\mathrm{i}\omega_+\,;1+\mathrm{i}\omega_1\,;\frac{1-\zeta}{2}\Big)\Big]\,, \end{aligned} \tag{B.19}$$

where

$$\tilde{\alpha} \equiv \Big(\frac{\omega_1}{\omega_0}\Big)^{1/2}\frac{\Gamma(1+\mathrm{i}\omega_0)\Gamma(\mathrm{i}\omega_1)}{\Gamma(1+\mathrm{i}\omega_+)\Gamma(\mathrm{i}\omega_+)}\ ,\ \ \tilde{\beta} \equiv -\Big(\frac{\omega_1}{\omega_0}\Big)^{1/2}\frac{\Gamma(1-\mathrm{i}\omega_0)\Gamma(\mathrm{i}\omega_1)}{\Gamma(1+\mathrm{i}\omega_-)\Gamma(\mathrm{i}\omega_-)}\,. \tag{B.20}$$

With these, the asymptotic forms for $t_1 \sim +\infty$ can be obtained easily as

$$f_1 = f(t_1) \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}\big(\tilde{\alpha}^* e^{-\mathrm{i}\omega_1 t_1} - \tilde{\beta} e^{\mathrm{i}\omega_1 t_1}\big)\,, \tag{B.21}$$

$$g_1 = g(t_1) \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}\big(\tilde{\alpha} e^{\mathrm{i}\omega_1 t_1} - \tilde{\beta}^* e^{-\mathrm{i}\omega_1 t_1}\big)\,, \tag{B.22}$$

and

$$u_1 \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}(-2\mathrm{i}\tilde{\beta}\omega_1)e^{\mathrm{i}\omega_1 t_1}\,, \quad \bar{u}_1 \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}(-2\mathrm{i}\tilde{\alpha}^*\omega_1)e^{-\mathrm{i}\omega_1 t_1}\,, \tag{B.23}$$

$$v_1 \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}(2\mathrm{i}\tilde{\alpha}\omega_1)e^{\mathrm{i}\omega_1 t_1}\,, \qquad \bar{v}_1 \sim \Big(\frac{\omega_0}{\omega_1}\Big)^{1/2}(2\mathrm{i}\tilde{\beta}^*\omega_1)e^{-\mathrm{i}\omega_1 t_1}\,. \tag{B.24}$$

We then find that the Bogoliubov coefficients take the asymptotic forms

$$\alpha(t_1,t_0) \sim \tilde{\alpha}\,e^{-\mathrm{i}(\omega_0 t_0 - \omega_1 t_1)} \equiv \alpha_1 \quad \big(\bar{\alpha}(t_1,t_0) \sim \alpha_1^*\big)\,, \tag{B.25}$$

$$\beta(t_1,t_0) \sim \tilde{\beta}\,e^{\mathrm{i}(\omega_0 t_0 + \omega_1 t_1)} \equiv \beta_1 \quad \big(\bar{\beta}(t_1,t_0) \sim \beta_1^*\big)\,. \tag{B.26}$$

They coincide with the well-known values in the literature (see, e.g., in [7]) up to a phase. It is easy to see

$$|\alpha_1|^2 = \frac{\sinh^2(\pi\omega_+)}{\sinh(\pi\omega_0)\sinh(\pi\omega_1)}\,, \quad |\beta_1|^2 = \frac{\sinh^2(\pi\omega_-)}{\sinh(\pi\omega_0)\sinh(\pi\omega_1)}\,, \tag{B.27}$$

and thus the relation $|\alpha_1|^2 - |\beta_1|^2 = 1$ actually holds.

Using the asymptotic forms of u_1 and v_1, we can calculate the wave function $\varphi(t, t_1)$ for $t_1 \to +\infty$:

$$\varphi(t; t_1) = \frac{1}{\sqrt{2\omega_0}} e^{\mathrm{i}\omega_1 t_1} \Big[\tilde{\alpha} f(t) + \tilde{\beta} g(t)\Big]\,. \tag{B.28}$$

This can be further rewritten by using Kummer's relation

$$F(a, b; c;\, z) = (1-z)^{c-a-b}\, F(c-a,\, c-b;\, c;\, z) \tag{B.29}$$

into the form

$$\varphi(t; t_1) = \frac{1}{\sqrt{2\omega_1}} e^{\mathrm{i}\omega_1 t_1 - \mathrm{i}\omega_+ t - \mathrm{i}\omega_- \log(2\cosh t)} F\Big(\mathrm{i}\omega_-,\, 1 + \mathrm{i}\omega_-\,; 1 - \mathrm{i}\omega_0\,; \frac{1-\tanh t}{2}\Big)\,. \tag{B.30}$$

This certainly coincides up to a phase with the positive-energy wave function in the remote future given in [7]. One can easily see that $\varphi(t; t_1)$ actually has the form $\varphi(t; t_1) \sim (1/\sqrt{2\omega_1})\, e^{-\mathrm{i}\omega_1 (t - t_1)}$ when t is also very large.

C. Propagator in Minkowski space

In order to evaluate the integral (3.28), we introduce the polar coordinates for the wave vector as

$$\mathrm{d}\boldsymbol{k}^2 = \mathrm{d}k^2 + k^2 \left(\mathrm{d}\theta^2 + \sin^2\theta\, \mathrm{d}\Omega_{d-3}^2\right), \tag{C.1}$$

where θ is chosen such that $\theta = 0$ corresponds to the direction $\boldsymbol{x} - \boldsymbol{x}'$, i.e., $\boldsymbol{k} \cdot (\boldsymbol{x} - \boldsymbol{x}') = k\, |\boldsymbol{x} - \boldsymbol{x}'| \cos\theta$. The volume element is then given by

$$\mathrm{d}^{d-1}\boldsymbol{k} = \mathrm{d}k\, k^{d-2}\, \mathrm{d}\theta\, \sin^{d-3}\theta\, \mathrm{d}\Omega_{d-3} = \mathrm{d}k\, k^{d-2}\, \mathrm{d}\cos\theta\, (1 - \cos^2\theta)^{(d-4)/2}\, \mathrm{d}\Omega_{d-3}\,. \tag{C.2}$$

Equation (3.28) then becomes

$$G(x, x') = \frac{|\Omega_{d-3}|}{(2\pi)^{d-1}} \int_0^\infty \mathrm{d}k\, k^{d-1}\, G_{\boldsymbol{k}}(t, t') \int_{-1}^1 \mathrm{d}s\, (1 - s^2)^{(d-4)/2} \cos\!\big(k|\boldsymbol{x} - \boldsymbol{x}'|\, s\big)\,, \tag{C.3}$$

where $|\Omega_{n-1}|$ is the area of the unit sphere in n-dimensional Euclidean space, $|\Omega_{n-1}| = \int \mathrm{d}\Omega_{n-1} = 2\,\pi^{n/2}/\Gamma(n/2)$. The integration with respect to $s = \cos\theta$ can be carried out by using the formula (8.411-8 in [22])

$$\int_{-1}^1 \mathrm{d}s\, (1 - s^2)^\nu \cos(zs) = \sqrt{\pi}\, \Gamma(\nu + 1) \left(\frac{z}{2}\right)^{-\nu - \frac{1}{2}} J_{\nu + \frac{1}{2}}(z) \quad [\,\mathrm{Re}\,\nu > -1\,]\,, \tag{C.4}$$

and we obtain

$$
\begin{aligned}
G(x,x') &= \frac{1}{(2\pi)^{\frac{d-1}{2}}\,|\boldsymbol{x}-\boldsymbol{x}'|^{\frac{d-3}{2}}}\int_0^\infty \mathrm{d}k\,k^{\frac{d-1}{2}}\,G_{\boldsymbol{k}}(t,t')\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big) \\
&= \frac{2^{-(d+1)/2}\,\pi^{-(d-1)/2}}{|\boldsymbol{x}-\boldsymbol{x}'|^{(d-3)/2}}\int_0^\infty \mathrm{d}k\,\frac{k^{(d-1)/2}}{\omega_k}\,e^{-\mathrm{i}\omega_k\,(t_>-t_<)}\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big)\,. \qquad \text{(C.5)}
\end{aligned}
$$

Although the integration does not converge in the UV region, this expression still has a meaning as a distribution and is analytic in the lower half of the complex $t_> - t_<$ plane. In order to make the integral manifestly converge, we introduce a small imaginary part by replacing $t_> - t_<$ by $t_> - t_< - \mathrm{i}0$ which does not alter the analytic property. Thus, what we need to evaluate is the integral

$$
\begin{aligned}
G(x,x') &= \frac{2^{-(d+1)/2}\,\pi^{-(d-1)/2}}{|\boldsymbol{x}-\boldsymbol{x}'|^{(d-3)/2}}\int_0^\infty \mathrm{d}k\,\frac{k^{(d-1)/2}}{\omega_k}\,e^{-e^{\mathrm{i}\pi/2}\,\omega_k\,(t_>-t_<-\mathrm{i}0)}\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big) \\
&= \frac{2^{-(d+1)/2}\,\pi^{-(d-1)/2}}{|\boldsymbol{x}-\boldsymbol{x}'|^{(d-3)/2}}\,m^{\frac{d-1}{2}}\int_1^\infty \mathrm{d}\lambda\,(\lambda^2-1)^{\frac{d-3}{4}}\,e^{-m\,\lambda\,e^{\mathrm{i}\pi/2}\,(t_>-t_<-\mathrm{i}0)}\,J_{\frac{d-3}{2}}\big(k|\boldsymbol{x}-\boldsymbol{x}'|\big)\,, \qquad \text{(C.6)}
\end{aligned}
$$

where, assuming $m > 0$, we have set $\lambda = \omega_k/m = \sqrt{k^2+m^2}/m$ to obtain the last expression. Then applying the formula (6.645-2 in [22]):

$$
\begin{aligned}
&\int_1^\infty \mathrm{d}\lambda\,(\lambda^2-1)^{\frac{\nu}{2}}\,e^{-\alpha\,\lambda}\,J_\nu\big(\beta\,\sqrt{\lambda^2-1}\big) \\
&\quad= \sqrt{\frac{2}{\pi}}\,\beta^\nu\,\big(\alpha^2+\beta^2\big)^{-\frac{\nu}{2}-\frac{1}{4}}\,K_{\nu+\frac{1}{2}}\big(\sqrt{\alpha^2+\beta^2}\big) \quad [\,\mathrm{Re}\,\alpha > 0\,,\ \beta\in\mathbb{R}\,] \qquad \text{(C.7)}
\end{aligned}
$$

with $\nu = (d-3)/2$, $\alpha = m\,e^{\mathrm{i}\pi/2}\,(t_> - t_< - \mathrm{i}0)$, $\beta = m\,|\boldsymbol{x}-\boldsymbol{x}'|$, we obtain

$$
G(x,x') = \frac{m^{(d-2)/2}}{(2\pi)^{d/2}}\,\big[e^{\mathrm{i}\pi}\,(\Delta t^2-\mathrm{i}0)+\Delta\boldsymbol{x}^2\big]^{-\frac{d-2}{4}}\,K_{\frac{d-2}{2}}\big(m\sqrt{e^{\mathrm{i}\pi}(\Delta t^2-\mathrm{i}0)+\Delta\boldsymbol{x}^2}\big)\,, \qquad \text{(C.8)}
$$

where $\Delta t \equiv t - t'$ and $\Delta\boldsymbol{x} \equiv \boldsymbol{x}-\boldsymbol{x}'$. This expression can be further rewritten by separately investigating the cases for different sign of $\sigma \equiv (x-x')^2 = -\Delta t^2 + \Delta\boldsymbol{x}^2$.

(1) spacelike $(\sigma > 0)$:

The modified Bessel function is readily evaluated as $K_{(d-2)/2}\big(m\sqrt{e^{\mathrm{i}\pi}(\Delta t^2-\mathrm{i}0)+\Delta\boldsymbol{x}^2}\big) = K_{(d-2)/2}\big(m\,\sqrt{\sigma}\big)$, and we have

$$
G(x,x') = \frac{m^{(d-2)/2}}{(2\pi)^{d/2}}\,\sigma^{-\frac{d-2}{4}}\,K_{\frac{d-2}{2}}\big(m\,\sqrt{\sigma}\big)\,. \qquad \text{(C.9)}
$$

(2) timelike $(\sigma < 0)$:

By using the relations $e^{\mathrm{i}\pi}\,(\Delta t^2-\mathrm{i}0)+\Delta\boldsymbol{x}^2 = e^{\mathrm{i}\pi}\,(-\sigma)$ and $K_\nu(e^{\mathrm{i}\pi/2}\,z) = -\,(\mathrm{i}\pi/2)\,H_\nu^{(2)}(z)$, we have

$$
G(x,x') = \frac{\pi}{2}\,\frac{m^{(d-2)/2}}{(2\pi)^{d/2}\,\mathrm{i}^{d-1}}\,(-\sigma)^{-\frac{d-2}{4}}\,H^{(2)}_{\frac{d-2}{2}}\big(m\,\sqrt{\sigma}\big)\,. \qquad \text{(C.10)}
$$

(3) null ($\sigma \to 0+$):

By using the expansion

$$K_\nu(z) = \frac{\Gamma(\nu)}{2}\left(\frac{z}{2}\right)^{-\nu}\big(1 + O(z^2)\big)\,, \tag{C.11}$$

we obtain

$$G(x,x') \to \frac{\Gamma\big((d-2)/2\big)}{4\,\pi^{d/2}}\,(\sigma + \mathrm{i}0)^{-\frac{d-2}{2}}\,. \tag{C.12}$$

The right-hand side actually coincides with the massless propagator.

It is easy to see that all of the expression for three cases can be derived from a single expression,

$$G(x,x') = \frac{m^{(d-2)/2}}{(2\pi)^{d/2}\,\big(\sigma + \mathrm{i}0\big)^{(d-2)/4}}\,K_{\frac{d-2}{2}}\big(m\sqrt{\sigma + \mathrm{i}0}\big)\,. \tag{C.13}$$

D. Proofs of Eqs. (4.47) and (4.51)

In order to show Eq. (4.47), we start from the following equation (6.578-10 in [22]):

$$\begin{aligned}
&\int_0^\infty \mathrm{d}x\, x^{\mu+1}\, K_\nu\big(\hat{a}\,x\big)\, K_\nu(\hat{b}\,x)\, J_\mu(c\,x) \\
&\qquad = \frac{\sqrt{\pi}\, c^\mu\, \Gamma(\mu+\nu+1)\,\Gamma(\mu-\nu+1)}{2^{3/2}\,(\hat{a}\,\hat{b})^{\mu+1}}\,\big(u^2-1\big)^{-\frac{1}{2}(\mu+\frac{1}{2})}\,\mathcal{P}_{\nu-1/2}^{-(\mu+\frac{1}{2})}(u) \\
&\Big[u \equiv \frac{\hat{a}^2 + \hat{b}^2 + c^2}{2\hat{a}\,\hat{b}}\,,\quad \mathrm{Re}\,(\hat{a}+\hat{b}) > |\mathrm{Im}\,c|\,,\quad \mathrm{Re}\,(\mu\pm\nu) > -1\,,\quad \mathrm{Re}\,\mu > -1\Big]\,.
\end{aligned} \tag{D.1}$$

By setting $\hat{a} = e^{-\mathrm{i}\pi/2}\,a$ and $\hat{b} = e^{\mathrm{i}\pi/2}\,b$ with $-\pi/2 < \arg a \le \pi$ and $-\pi/2 < \arg b \le \pi$, and by using the identities

$$H_\nu^{(1)}(a\,x) = -\frac{2\mathrm{i}}{\pi}\,e^{-\mathrm{i}\pi\nu/2}\,K_\nu\big(\hat{a}\,x\big)\,,\quad H_\nu^{(2)}(b\,x) = \frac{2\mathrm{i}}{\pi}\,e^{\mathrm{i}\pi\nu/2}\,K_\nu\big(\hat{b}\,x\big)\,, \tag{D.2}$$

the equation is rewritten to the form

$$\begin{aligned}
&\int_0^\infty \mathrm{d}x\, x^{\mu+1}\, H_\nu^{(1)}(a\,x)\, H_\nu^{(2)}(b\,x)\, J_\mu(c\,x) \\
&\qquad = \frac{\sqrt{2}\, c^\mu\, \Gamma(\mu+\nu+1)\,\Gamma(\mu-\nu+1)}{\pi^{3/2}\,(\hat{a}\,\hat{b})^{\mu+1}}\,\big(u^2-1\big)^{-\frac{1}{2}(\mu+\frac{1}{2})}\,\mathcal{P}_{\nu-1/2}^{-\mu-\frac{1}{2}}(u) \\
&\Big[u \equiv \frac{-a^2 - b^2 + c^2}{2a\,b}\,,\quad \mathrm{Im}\,(a-b) > |\mathrm{Im}\,c|\,,\quad \mathrm{Re}\,(\mu\pm\nu) > -1\,,\quad \mathrm{Re}\,\mu > -1\Big]\,.
\end{aligned} \tag{D.3}$$

We substitute to this $x = k$, $a = -\eta_< - \mathrm{i}\varepsilon'$, $b = -\eta_> - \mathrm{i}\varepsilon$, $c = |\boldsymbol{x} - \boldsymbol{x}'|$ and $\mu = (d-3)/2$ where the convergence of the integral requires the inequality $\varepsilon > \varepsilon'$. We then obtain Eq. (4.47) with $u = -Z(x,x') + \mathrm{i}\,0$ for infinitesimal ε and ε'.

In order to show Eq. (4.51), we use the following equation (6.578-11 in [22]):

$$\int_0^\infty \mathrm{d}x\, x^{\mu+1}\, K_\nu\big(\hat{a}\,x\big)\, I_\nu\big(\hat{b}\,x\big)\, J_\mu(c\,x) = \frac{c^\mu\, e^{-\mathrm{i}\pi(\mu+\frac{1}{2})}}{\sqrt{2\pi}\,(\hat{a}\,\hat{b})^{\mu+1}}\, \big(u^2-1\big)^{-\frac{1}{2}(\mu+\frac{1}{2})}\, \mathcal{Q}^{\mu+\frac{1}{2}}_{\nu-1/2}(u)$$

$$\bigg[u \equiv \frac{\hat{a}^2+\hat{b}^2+c^2}{2\hat{a}\,\hat{b}}\,, \quad \mathrm{Re}\,\hat{a} > |\mathrm{Re}\,\hat{b}| + |\mathrm{Im}\,c|\,, \quad \mathrm{Re}\,\mu > -1\,, \quad \mathrm{Re}\,(\mu+\nu) > -1\bigg]\,. \tag{D.4}$$

By setting $\hat{a} = e^{\mathrm{i}\pi/2}\,a$ and $\hat{b} = e^{\mathrm{i}\pi/2}\,b$ for $-\pi/2 < \arg a \le \pi$ and $-\pi < \arg b \le \pi/2$, and by using the identities

$$H^{(2)}_\nu(a\,x) = \frac{2\mathrm{i}}{\pi}\, e^{\mathrm{i}\pi\nu/2}\, K_\nu\big(\hat{a}\,x\big)\,, \quad J_\nu(b\,x) = e^{-\mathrm{i}\pi\nu/2}\, I_\nu\big(\hat{b}\,x\big)\,, \tag{D.5}$$

the equation is rewritten to the form

$$\int_0^\infty \mathrm{d}x\, x^{\mu+1}\, H^{(2)}_\nu(a\,x)\, J_\nu(b\,x)\, J_\mu(c\,x) = \frac{\sqrt{2}\,c^\mu\, e^{-2\mathrm{i}\pi(\mu+\frac{1}{2})}}{\pi^{3/2}\,(a\,b)^{\mu+1}}\, \big(u^2-1\big)^{-\frac{1}{2}(\mu+\frac{1}{2})}\, \mathcal{Q}^{\mu+\frac{1}{2}}_{\nu-1/2}(u)$$

$$\bigg[u \equiv \frac{a^2+b^2-c^2}{2a\,b}\,, \quad (-\mathrm{Im}\,a) > |\mathrm{Im}\,b| + |\mathrm{Im}\,c|\,, \quad \mathrm{Re}\,\mu > -1\,, \quad \mathrm{Re}\,(\mu+\nu) > -1\bigg]\,. \tag{D.6}$$

We substitute to this $x = k$, $a = -\eta_< - \mathrm{i}\varepsilon$, $b = -\eta_> - \mathrm{i}\varepsilon'$, $c = |\boldsymbol{x} - \boldsymbol{x}'|$ and $\mu = (d-3)/2$, where the convergence of the integral requires the inequality $\varepsilon > \varepsilon'$. We then obtain Eq. (4.51) with $u = Z(x,x') - \mathrm{i}\,0$ for infinitesimal ε and ε'.

E. Associated Legendre functions and the addition formulae

In this appendix, we give several formulae of the associated Legendre functions which are used in subsec. 4.3. For details of the associated Legendre functions, see [22] and [23].

The associated Legendre functions $\mathcal{P}^\mu_\nu(z)$ and $\mathcal{Q}^\mu_\nu(z)$ are defined over the complex z-plane other than the cut along the real axis to the left of the point $z = 1$ (running from $-\infty$ to 1), while the associated Legendre functions $\mathsf{P}^\mu_\nu(x)$ and $\mathsf{Q}^\mu_\nu(x)$ are defined only on the interval

$-1 < x < 1$:

$$\mathcal{P}_\nu^\mu(z) \equiv \frac{1}{\Gamma(1-\mu)} \left(\frac{z+1}{z-1}\right)^{\frac{\mu}{2}} F\Big(-\nu,\, \nu+1;\, 1-\mu;\, \frac{1-z}{2}\Big)\,, \tag{E.1}$$

$$\mathcal{Q}_\nu^\mu(z) \equiv \frac{e^{\mathrm{i}\pi\mu}\, \pi^{\frac{1}{2}}\, \Gamma(\nu+\mu+1)}{2^{\nu+1}\, \Gamma(\nu+3/2)}\, z^{-\nu-\mu-1} \left(z^2-1\right)^{\frac{\mu}{2}} F\Big(\frac{\nu+\mu+2}{2},\, \frac{\nu+\mu+1}{2};\, \nu+\frac{3}{2};\, \frac{1}{z^2}\Big)\,, \tag{E.2}$$

$$\begin{aligned} \mathsf{P}_\nu^\mu(x) &\equiv \frac{1}{2}\left[e^{\frac{1}{2}\,\mathrm{i}\pi\mu}\, \mathcal{P}_\nu^\mu(x+\mathrm{i}0) + e^{-\frac{1}{2}\,\mathrm{i}\pi\mu}\, \mathcal{P}_\nu^\mu(x-\mathrm{i}0)\right] \\ &= \frac{1}{\Gamma(1-\mu)} \left(\frac{1+x}{1-x}\right)^{\frac{\mu}{2}} F\Big(-\nu,\, \nu+1;\, 1-\mu;\, \frac{1-x}{2}\Big)\,, \end{aligned} \tag{E.3}$$

$$\begin{aligned} \mathsf{Q}_\nu^\mu(x) &\equiv \frac{1}{2}\, e^{-\mathrm{i}\pi\mu} \left[e^{-\frac{1}{2}\,\mathrm{i}\pi\mu}\, \mathcal{Q}_\nu^\mu(x+\mathrm{i}0) + e^{\frac{1}{2}\,\mathrm{i}\pi\mu}\, \mathcal{Q}_\nu^\mu(x-\mathrm{i}0)\right] \\ &= \frac{\pi}{2\sin\pi\mu}\left[\cos\pi\mu\, \mathsf{P}_\nu^\mu(x) - \frac{\Gamma(\nu+\mu+1)}{\Gamma(\nu-\mu+1)}\, \mathsf{P}_\nu^{-\mu}(x)\right]. \end{aligned} \tag{E.4}$$

functional relations

The four functions $\mathcal{P}(z)$, $\mathcal{Q}(z)$, $\mathsf{P}(x)$ and $\mathsf{Q}(x)$ are related to each other as (3.4 and 3.3.1 in [23])

$$e^{\mathrm{i}\pi\frac{\mu}{2}}\, \mathcal{P}_\nu^\mu(x+\mathrm{i}0) = e^{-\mathrm{i}\pi\frac{\mu}{2}}\, \mathcal{P}_\nu^\mu(x-\mathrm{i}0) = \mathsf{P}_\nu^\mu(x)\,, \tag{E.5}$$

$$e^{-\mathrm{i}\pi\mu}\left[e^{-\mathrm{i}\pi\frac{\mu}{2}}\, \mathcal{Q}_\nu^\mu(x+\mathrm{i}0) \pm e^{\mathrm{i}\pi\frac{\mu}{2}}\, \mathcal{Q}_\nu^\mu(x-\mathrm{i}0)\right] = \left\{\begin{array}{l} 2\mathsf{Q}_\nu^\mu(x) \\ -\mathrm{i}\pi\mathsf{P}_\nu^\mu(x) \end{array}\right.\,, \tag{E.6}$$

$$\mathcal{Q}_\nu^\mu(-z) = \left\{\begin{array}{ll} -\,e^{\mathrm{i}\pi\nu}\mathcal{Q}_\nu^\mu(z) & [\mathrm{Im}\, z > 0] \\ -\,e^{-\mathrm{i}\pi\nu}\mathcal{Q}_\nu^\mu(z) & [\mathrm{Im}\, z < 0] \end{array}\right.\,. \tag{E.7}$$

We also have (8.73 in [22])

$$\mathcal{P}_\nu^{-\mu}(z) = \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)}\left[\mathcal{P}_\nu^\mu(z) - \frac{2}{\pi}\, e^{-\mathrm{i}\pi\mu}\, \sin\pi\mu\, \mathcal{Q}_\nu^\mu(z)\right], \tag{E.8}$$

$$\mathcal{Q}_\nu^{-\mu}(z) = e^{-2\mathrm{i}\pi\mu}\, \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)}\, \mathcal{Q}_\nu^\mu(z)\,, \tag{E.9}$$

$$\mathsf{P}_\nu^{-\mu}(x) = \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)}\left[\cos\pi\mu\, \mathsf{P}_\nu^\mu(x) - \frac{2}{\pi}\, \sin\pi\mu\, \mathsf{Q}_\nu^\mu(x)\right], \tag{E.10}$$

$$\mathsf{Q}_\nu^{-\mu}(x) = \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)}\left[\frac{\pi}{2}\, \sin\pi\mu\, \mathsf{P}_\nu^\mu(x) + \cos\pi\mu\, \mathsf{Q}_\nu^\mu(x)\right], \tag{E.11}$$

$$\mathsf{P}_\nu^{-\mu}(-x) = \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)}\left[\cos\pi\nu\, \mathsf{P}_\nu^\mu(x) - \frac{2}{\pi}\, \sin\pi\nu\, \mathsf{Q}_\nu^\mu(x)\right], \tag{E.12}$$

$$\mathsf{P}_{-\nu-1}^\mu(x) = \mathsf{P}_\nu^\mu(x)\,, \tag{E.13}$$

$$\mathsf{Q}_{-\nu-1}^\mu(x) = \frac{1}{\sin\pi(\nu-\mu)}\big[-\pi\cos\pi\nu\, \mathsf{P}_\nu^\mu(x) + \sin\pi(\nu+\mu)\, \mathsf{Q}_\nu^\mu(x)\big]\,. \tag{E.14}$$

Their Wronskians have the forms (8.741 in [22])

$$\begin{vmatrix} \mathsf{P}^{\mu}_{\nu}(x) & \mathsf{Q}^{\mu}_{\nu}(x) \\ \frac{\mathrm{d}}{\mathrm{d}x}\mathsf{P}^{\mu}_{\nu}(x) & \frac{\mathrm{d}}{\mathrm{d}x}\mathsf{Q}^{\mu}_{\nu}(x) \end{vmatrix} = \frac{1}{1-x^2}\,\frac{\Gamma(\nu+\mu+1)}{\Gamma(\nu-\mu+1)}\,, \tag{E.15}$$

$$\begin{vmatrix} \mathsf{P}^{-\mu}_{\nu}(x) & \mathsf{P}^{\mu}_{\nu}(x) \\ \frac{\mathrm{d}}{\mathrm{d}x}\mathsf{P}^{-\mu}_{\nu}(x) & \frac{\mathrm{d}}{\mathrm{d}x}\mathsf{P}^{\mu}_{\nu}(x) \end{vmatrix} = \frac{1}{1-x^2}\,\frac{2\sin\pi\mu}{\pi}\,. \tag{E.16}$$

We also have (8.733-1 in [22])

$$(1-x^2)\,\frac{\mathrm{d}}{\mathrm{d}x}\,\mathsf{P}^{\mu}_{\nu}(x) = (\nu+1)\,x\,\mathsf{P}^{\mu}_{\nu}(x) - (\nu-\mu+1)\,\mathsf{P}^{\mu}_{\nu+1}(x)\,, \tag{E.17}$$

$$(1-x^2)\,\frac{\mathrm{d}}{\mathrm{d}x}\,\mathsf{Q}^{\mu}_{\nu}(x) = (\nu+1)\,x\,\mathsf{Q}^{\mu}_{\nu}(x) - (\nu-\mu+1)\,\mathsf{Q}^{\mu}_{\nu+1}(x)\,. \tag{E.18}$$

asymptotic forms

The associated Legendre functions have the following asymptotic forms near boundaries (3.9.2 in [23]):

$$\mathcal{P}^{\mu}_{\nu}(z) \overset{z\sim\infty}{\sim} \begin{cases} 2^{\nu}\,\pi^{-\frac{1}{2}}\,\dfrac{\Gamma(\nu+1/2)}{\Gamma(\nu-\mu+1)}\,z^{\nu} & [\,\mathrm{Re}\,\nu > -1/2\,] \\ 2^{-\nu-1}\,\pi^{-\frac{1}{2}}\,\dfrac{\Gamma(-\nu-1/2)}{\Gamma(-\nu-\mu)}\,z^{-\nu-1} & [\,\mathrm{Re}\,\nu < -1/2\,] \end{cases}\,, \tag{E.19}$$

$$\mathcal{Q}^{\mu}_{\nu}(z) \overset{z\sim\infty}{\sim} e^{\mathrm{i}\pi\mu}\,2^{-\nu-1}\,\pi^{\frac{1}{2}}\,\frac{\Gamma(\nu+\mu+1)}{\Gamma(\nu+3/2)}\,z^{-\nu-1}\,, \tag{E.20}$$

$$\mathsf{P}^{\nu}_{k}(x) \overset{x\sim+1}{\sim} \frac{2^{\nu}}{\Gamma(1-\nu)}\,\bigl(1-x^2\bigr)^{-\frac{\nu}{2}}\,, \tag{E.21}$$

$$\mathsf{Q}^{\nu}_{k}(x) \overset{x\sim+1}{\sim} 2^{\nu-1}\,\cos(\pi\nu)\,\Gamma(\nu)\,\bigl(1-x^2\bigr)^{-\frac{\nu}{2}} \qquad [\,\mathrm{Re}\,\nu > 0\,]\,, \tag{E.22}$$

$$\mathsf{P}^{\nu}_{k}(x) \overset{x\sim-1}{\sim} \begin{cases} 2^{-\nu}\,\cos(\pi k)\,\dfrac{\Gamma(k+\nu+1)}{\Gamma(\nu+1)\,\Gamma(k-\nu+1)}\bigl(1-x^2\bigr)^{\frac{\nu}{2}} & [\,k\in\mathbb{Z}\,] \\ -\,\dfrac{2^{\nu}\,\sin(\pi k)\,\Gamma(\nu)}{\pi}\,\bigl(1-x^2\bigr)^{-\frac{\nu}{2}} & [\,k\in\mathbb{Z}+1/2\,,\ \mathrm{Re}\,\nu>0\,] \end{cases}\,, \tag{E.23}$$

$$\mathsf{Q}^{\nu}_{k}(x) \overset{x\sim-1}{\sim} \begin{cases} -\,2^{\nu-1}\cos(\pi k)\,\Gamma(\nu)\,\bigl(1-x^2\bigr)^{-\frac{\nu}{2}} & [\,k\in\mathbb{Z}\,,\ \mathrm{Re}\,\nu>0\,] \\ -\,\dfrac{2^{-\nu-1}\,\pi\,\Gamma(k+\nu+1)}{\sin(\pi k)\,\Gamma(\nu+1)\,\Gamma(k-\nu+1)}\bigl(1-x^2\bigr)^{\frac{\nu}{2}} & [\,k\in\mathbb{Z}+1/2\,] \end{cases}\,. \tag{E.24}$$

addition formulae

We find the following formulae, which are useful in obtaining the propagators in the global patch:

$$\begin{aligned}&\frac{\mathrm{i}\pi(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}}{2\,(d-2)\,|\Omega_{d-1}|\,\sin(\pi\nu)}\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathsf{P}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\\&\quad=\frac{-\,e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\sin(\pi\nu)}\Big[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_+)-(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_-)\Big]\quad(d:\text{odd})\,,\end{aligned}\tag{E.25}$$

$$\begin{aligned}&\frac{\mathrm{i}(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}}{(d-2)\,|\Omega_{d-1}|\,\cos(\pi\nu)}\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathsf{Q}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\\&\quad=\frac{\mathrm{i}e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}\,\cos(\pi\nu)}\Big[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_+)+(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_-)\Big]\quad(d:\text{even})\,,\end{aligned}\tag{E.26}$$

$$\begin{aligned}&\frac{\mathrm{i}(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}}{(d-2)\,|\Omega_{d-1}|}\sum_{L=0}^{\infty}(2L+d-2)\,\frac{\Gamma(k-\nu+1)}{\Gamma(k+\nu+1)}\,\mathsf{P}_k^{\nu}(\sin\varphi_1)\,\mathsf{Q}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\\&\quad=\frac{\mathrm{i}e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}}\Big[e^{-\mathrm{i}\pi\nu}\,(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{-\nu-\frac12}^{\frac{d-2}{2}}(u_+)+e^{\mathrm{i}\pi\nu}\,(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{-\nu-\frac12}^{\frac{d-2}{2}}(u_-)\Big]\\&\qquad(d:\text{even})\,,\end{aligned}\tag{E.27}$$

$$\begin{aligned}&\frac{2\,\mathrm{i}(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}}{\pi\,(d-2)\,|\Omega_{d-1}|}\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{Q}_k^{-\nu}(\sin\varphi_1)\,\mathsf{Q}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\\&\quad=\frac{e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}}\Big\{(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_+)-(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_-)\\&\qquad\qquad+\frac{\mathrm{i}\,\pi}{\cos(\pi\nu)}\big[e^{-\mathrm{i}\pi\nu}\,(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}_{\nu-\frac12}^{\frac{d-2}{2}}(u_+)+e^{\mathrm{i}\pi\nu}\,(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}_{\nu-\frac12}^{\frac{d-2}{2}}(u_-)\big]\Big\}\\&\quad=-\frac{e^{-\mathrm{i}\pi\frac{d-2}{2}}}{2\,(2\pi)^{\frac{d}{2}}}\Big\{(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_+)-(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{Q}_{\nu-\frac12}^{\frac{d-2}{2}}(u_-)\\&\qquad\qquad-\frac{\pi}{\cos(\pi\nu)}\big[(u_+^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}_{\nu-\frac12}^{\frac{d-2}{2}}(-u_+)-(u_-^2-1)^{-\frac{d-2}{4}}\,\mathcal{P}_{\nu-\frac12}^{\frac{d-2}{2}}(-u_-)\big]\Big\}\\&\qquad(d:\text{even})\,,\end{aligned}\tag{E.28}$$

where $-\pi/2<\varphi_2<\varphi_1<\pi/2\,,\ 0\le\theta\le\pi\,,\ k\equiv L+(d-3)/2\,,$ and

$$u_\pm(\varphi_1,\varphi_2,\theta)\equiv-Z(\varphi_1,\varphi_2,\theta)\pm\mathrm{i}0\ \ \text{with}\ \ Z(\varphi_1,\varphi_2,\theta)\equiv\frac{-\sin\varphi_1\,\sin\varphi_2+\cos\theta}{\cos\varphi_1\,\cos\varphi_2}\,.\tag{E.29}$$

We prove Eqs. (E.25) and (E.26) for the rest of this appendix. Equation (E.27) can be proved in a similar way, and (E.28) is readily obtained from (E.26) and (E.27).

We start from Eq. (12) of [24]:[21]

$$\begin{aligned}
&(\sinh\gamma)^{-\frac{d-2}{2}}\,\mathcal{Q}_{\nu-\frac{1}{2}}^{\frac{d-2}{2}}(\cosh\gamma)\\
&\qquad= 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\left(-\nu+\frac{d-2}{2}\right)}\,(\sinh\beta_1\,\sinh\beta_2)^{\frac{d-1}{2}}\\
&\qquad\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathcal{P}_k^{-\nu}(\cosh\beta_1)\,\mathcal{Q}_k^{\nu}(\cosh\beta_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\,,
\end{aligned}\tag{E.30}$$

where

$$\cosh\gamma(\beta_1,\beta_2,\theta)\equiv\frac{\cosh\beta_1\cosh\beta_2-\cos\alpha}{\sinh\beta_1\sinh\beta_2}\,,\quad \mathrm{Re}\,\beta_2>|\mathrm{Re}\,\beta_1|+|\mathrm{Im}\,\theta|\,.\tag{E.31}$$

Both sides of Eq. (E.30) should be understood as the quantities that are continued analytically from the region where β_1, β_2, and θ take all real values (for which $\cosh\gamma>1$). We reparametrize the variables in Eq. (E.30) as

$$\begin{gathered}
\beta_\alpha^{\pm}\equiv\pm\mathrm{i}\Big(\frac{\pi}{2}-\varphi_\alpha\Big)+\varepsilon_\alpha\quad(\alpha=1,2)\\
\Big[-\frac{\pi}{2}<\varphi_2<\varphi_1<\frac{\pi}{2}\,,\quad 0<\varepsilon_1<\varepsilon_2\ll 1\Big]\,,
\end{gathered}\tag{E.32}$$

and only keep the contributions from ε_α to the linear order. We then have

$$\cosh\beta_\alpha^{\pm}=\sin\varphi_\alpha\pm\mathrm{i}\varepsilon_\alpha\cos\phi_\alpha\,,\qquad \sinh\beta_\alpha^{\pm}=\pm\mathrm{i}\cos\varphi_\alpha+\varepsilon_\alpha\sin\phi_\alpha\,,\tag{E.33}$$

and

$$\cosh\gamma_\pm\equiv\cosh\gamma(\beta_1^{\mp},\beta_2^{\pm},\theta)=-Z(\varphi_1,\varphi_2,\theta)+\mathrm{i}\mathcal{O}(\varepsilon_1,\varepsilon_2)\,.\tag{E.34}$$

If we fix the parameters ε_1 and ε_2, and vary φ_1, φ_2, and θ within the regions $-\pi/2<\varphi_2<\varphi_1<\pi/2$ and $0\le\theta\le\pi$, then $\cosh\gamma_\pm$ ranges in the region depicted in Fig. 6.

In the following, we divide the parameter region of φ_1, φ_2 and θ into three parts, where Z takes values in (1) $Z>1$, (2) $1>Z>-1$, and (3) $Z<-1$, respectively. We then derive a simpler expression of Eq. (E.30) for each case, and show that the obtained expressions for the three cases can be summarized in the form (E.25) and (E.26).

(1) $Z>1$

In this case, $\cosh\gamma_\pm=-Z\pm\mathrm{i}0$ as can be seen from Fig. 6.[22] Thus, we have

$$\sinh\gamma_\pm=e^{\pm\mathrm{i}\pi}\left(Z^2-1\right)^{\frac{1}{2}}\,,\tag{E.35}$$

$$\mathcal{Q}_{\nu-\frac{1}{2}}^{\frac{d-2}{2}}(\cosh\gamma_\pm)=-\,e^{\mp\mathrm{i}\pi(\nu-\frac{1}{2})}\,\mathcal{Q}_{\nu-\frac{1}{2}}^{\frac{d-2}{2}}(Z\mp\mathrm{i}0)=-\,e^{\mp\mathrm{i}\pi(\nu-\frac{1}{2})}\,\mathcal{Q}_{\nu-\frac{1}{2}}^{\frac{d-2}{2}}(Z)\,,\tag{E.36}$$

---

[21] Equation (E.27) can be proved by replacing $\mathcal{Q}_k^{\nu}(\cosh\beta_2)$ in (E.30) by $\mathcal{Q}_k^{-\nu}(\cosh\beta_2)$ with the help of (E.9).

[22] ε_1 and ε_2 are to be taken to zero with keeping $\varepsilon_2>\varepsilon_1$.



Figure 6: Schematic view illustrating the range of $\cosh\gamma_+$ for $\varepsilon_1 = 0.001$ and $\varepsilon_2 = 0.02$. Here, φ_1 and φ_2 run over the range $-\pi/2 < \varphi_2 < \varphi_1 < \pi/2$, and θ runs over its full range $0 \leq \theta \leq \pi$. The range of $\cosh\gamma_-$ can be obtained by turning the figure by 180° over the horizontal axis.

where we have used Eq. (E.7) and the fact that $\mathcal{Q}^{\mu}_{\nu}(z)$ does not have a cut in the region $\mathrm{Re}\, z > 1$. Then, Eq. (E.30) becomes

$$\begin{aligned}
&- e^{\mp \mathrm{i}\pi(\frac{d-3}{2}+\nu)} \left(Z^2-1\right)^{-\frac{d-2}{4}} \mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(Z) \\
&= 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\, e^{\mathrm{i}\pi\left(-\nu\mp\frac{\nu}{2}+\frac{d-2}{2}\right)} \left(\cos\varphi_1\,\cos\varphi_2\right)^{\frac{d-1}{2}} \\
&\quad\times \sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\,\mathcal{Q}^{\nu}_{k}(\sin\varphi_2 \pm \mathrm{i}0)\, C^{\frac{d-2}{2}}_{L}(\cos\theta)\,. \qquad \text{(E.37)}
\end{aligned}$$

By taking the difference between the above equations with the upper and the lower signs, we obtain

$$\begin{aligned}
&2\mathrm{i}\sin\Big[\pi\Big(\frac{d-3}{2}+\nu\Big)\Big] \left(Z^2-1\right)^{-\frac{d-2}{4}} \mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(Z) \\
&= -\,\mathrm{i}\pi 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\, e^{\mathrm{i}\pi\frac{d-2}{2}} \left(\cos\varphi_1\,\cos\varphi_2\right)^{\frac{d-1}{2}} \\
&\quad\times \sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\,\mathsf{P}^{\nu}_{k}(\sin\varphi_2)\, C^{\frac{d-2}{2}}_{L}(\cos\theta)\,, \qquad \text{(E.38)}
\end{aligned}$$

where Eq. (E.6) has been used. Similarly, by taking their sum, we obtain

$$\begin{aligned}
&-2\cos\Big[\pi\Big(\frac{d-3}{2}+\nu\Big)\Big]\,\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(Z) \\
&= 2^{\frac{d}{2}-1}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathsf{Q}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\,. \qquad \text{(E.39)}
\end{aligned}$$

The right hand side of Eqs. (E.25) and (E.26) can then be written as

$$\begin{cases}
-\dfrac{\mathrm{i}\sin\pi(\frac{d-2}{2})}{(2\pi)^{\frac{d}{2}}}\,e^{-\mathrm{i}\pi\frac{d-2}{2}}\,\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(Z) & (d:\text{odd})\,, \\
-\dfrac{\mathrm{i}\cos\pi(\frac{d-2}{2})\tan\pi\nu}{(2\pi)^{\frac{d}{2}}}\,e^{-\mathrm{i}\pi\frac{d-2}{2}}\,\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(Z) & (d:\text{even})\,.
\end{cases} \qquad \text{(E.40)}$$

(2) $1 > Z > -1$

In this case, as can be seen from Fig. 6, $\cosh\gamma_+$ crosses the branch cut between $-1 < z < 1$ from above and move to another Riemann sheet. On the other hand, $\cosh\gamma_-$ crosses the branch cut between $-1 < z < 1$ from below. Thus, in this region, we have

$$\sinh\gamma_\pm = e^{\pm\mathrm{i}\frac{\pi}{2}}\,\big(1-Z^2\big)^{\frac{1}{2}}\,, \qquad \text{(E.41)}$$

$$\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(\cosh\gamma_\pm) = \mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z\pm\mathrm{i}0) = e^{\mathrm{i}\pi\frac{d-2}{2}(1\pm\frac{1}{2})}\Big[\mathsf{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\mp\frac{\mathrm{i}\pi}{2}\,\mathsf{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\Big]\,, \qquad \text{(E.42)}$$

where we have used Eq. (E.6). Then, Eq. (E.30) becomes

$$\begin{aligned}
&e^{\mathrm{i}\pi\frac{d-2}{2}}\,\big(1-Z^2\big)^{-\frac{d-2}{4}}\Big[\mathsf{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\mp\frac{\mathrm{i}\pi}{2}\,\mathsf{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\Big] \\
&= 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\left(-\nu\mp\frac{\nu}{2}+\frac{d-2}{2}\right)}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathcal{Q}_k^{\nu}(\sin\varphi_2\pm\mathrm{i}0)\,C_L^{\frac{d-2}{2}}(\cos\theta)\,. \qquad \text{(E.43)}
\end{aligned}$$

By taking the difference of the equations with the upper and the lower signs, we obtain

$$\begin{aligned}
\big(1-Z^2\big)^{-\frac{d-2}{4}}\,\mathsf{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) &= 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathsf{P}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\,. \qquad \text{(E.44)}
\end{aligned}$$

Similarly, by taking their sum, we obtain

$$\begin{aligned}
\big(1-Z^2\big)^{-\frac{d-2}{4}}\,\mathsf{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) &= 2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}_k^{-\nu}(\sin\varphi_1)\,\mathsf{P}_k^{\nu}(\sin\varphi_2)\,C_L^{\frac{d-2}{2}}(\cos\theta)\,. \qquad \text{(E.45)}
\end{aligned}$$

The right hand side of Eqs. (E.25) and (E.26) can then be written as follows:

$$\begin{cases} \dfrac{\mathrm{i}\pi}{2\,(2\pi)^{\frac{d}{2}}\sin(\pi\nu)}\,\big(1-Z^2\big)^{-\frac{d-2}{4}}\,\mathsf{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) & (d:\mathrm{odd})\,, \\ \dfrac{\mathrm{i}}{(2\pi)^{\frac{d}{2}}\cos(\pi\nu)}\,\big(1-Z^2\big)^{-\frac{d-2}{4}}\,\mathsf{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) & (d:\mathrm{even})\,. \end{cases} \tag{E.46}$$

(3) $Z<-1$

In this case, from Fig. 6, we know that, in the region $\mathrm{Re}\,z>1$, $\cosh\gamma_+$ runs above the real axis, or below the real axis in the next Riemann sheet after passing through the cut on $-1<z<1$ from above. On the other hand, in the region $\mathrm{Re}\,z>1$, $\cosh\gamma_-$ runs below the real axis, or above the real axis in another sheet after passing through the cut on $-1<z<1$ from below.

When $\cosh\gamma_\pm=-Z\pm\mathrm{i}0$, we have

$$\sinh\gamma_\pm=\big(Z^2-1\big)^{\frac{1}{2}}\,,\qquad \mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(\cosh\gamma_\pm)=\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\,, \tag{E.47}$$

and then the following equation is obtained:

$$\begin{aligned} &\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) \\ &=2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\left(-\nu\mp\frac{\nu}{2}+\frac{d-2}{2}\right)}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\ &\qquad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_k(\sin\varphi_1)\,\mathcal{Q}^{\nu}_k(\sin\varphi_2\pm\mathrm{i}0)\,C^{\frac{d-2}{2}}_L(\cos\theta)\,. \end{aligned} \tag{E.48}$$

By taking the difference and the sum of the above equations with the upper and the lower signs, we obtain

$$\begin{aligned} 0=&\,-\mathrm{i}\pi\,2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\ &\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_k(\sin\varphi_1)\,\mathsf{P}^{\nu}_k(\sin\varphi_2)\,C^{\frac{d-2}{2}}_L(\cos\theta)\,, \end{aligned} \tag{E.49}$$

$$\begin{aligned} 2\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)=&\,2^{\frac{d}{2}-1}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}} \\ &\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_k(\sin\varphi_1)\,\mathsf{Q}^{\nu}_k(\sin\varphi_2)\,C^{\frac{d-2}{2}}_L(\cos\theta)\,. \end{aligned} \tag{E.50}$$

On the other hand, when $\cosh\gamma_\pm=-Z\mp\mathrm{i}0$, we need to evaluate the functions $\sinh\gamma_\pm$ and $\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(\cosh\gamma_\pm)$ on a new Riemann sheet, since $\cosh\gamma_\pm$ has already crossed the branch cut. We then have

$$\sinh\gamma_\pm=e^{\pm\mathrm{i}\pi}\,\big(Z^2-1\big)^{\frac{1}{2}}\,, \tag{E.51}$$

$$\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(\cosh\gamma_\pm)=e^{\pm\mathrm{i}\pi\frac{d-2}{2}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\mp\mathrm{i}\pi\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\,, \tag{E.52}$$

and Eq. (E.30) takes the form

$$\begin{aligned}
&\big(Z^2-1\big)^{-\frac{d-2}{4}}\Big[\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\mp \mathrm{i}\pi\, e^{\pm\mathrm{i}\pi\frac{d-2}{2}(1\mp 1)}\,\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\Big]\\
&=2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\left(-\nu\mp\frac{\nu}{2}+\frac{d-2}{2}\right)}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}\\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\,\mathcal{Q}^{\nu}_{k}(\sin\varphi_2\pm\mathrm{i}0)\,C^{\frac{d-2}{2}}_{L}(\cos\theta)\,. \qquad \text{(E.53)}
\end{aligned}$$

By taking the difference of the above equations with the upper and the lower signs, we obtain

$$\begin{aligned}
&\cos\Big(\frac{d-2}{2}\pi\Big)\big(Z^2-1\big)^{-\frac{d-2}{4}}\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\\
&=2^{\frac{d}{2}-2}\Gamma\Big(\frac{d-2}{2}\Big)e^{\mathrm{i}\pi\frac{d-2}{2}}(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}\sum_{L=0}^{\infty}(2L+d-2)\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\mathsf{P}^{\nu}_{k}(\sin\varphi_2)C^{\frac{d-2}{2}}_{L}(\cos\theta)\,.
\end{aligned} \qquad \text{(E.54)}$$

In particular, in odd dimensions, we have

$$\begin{aligned}
0&=2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}\\
&\quad\times\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\,\mathsf{P}^{\nu}_{k}(\sin\varphi_2)\,C^{\frac{d-2}{2}}_{L}(\cos\theta)\,, \qquad \text{(E.55)}
\end{aligned}$$

which is equivalent to Eq. (E.49). Similarly, by taking the sum of Eq. (E.53) with the upper and the lower signs, we have

$$\begin{aligned}
&\big(Z^2-1\big)^{-\frac{d-2}{4}}\Big[\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)-\pi e^{\mathrm{i}\pi\frac{d-2}{2}}\sin\Big(\frac{d-2}{2}\pi\Big)\mathcal{P}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\Big]\\
&=2^{\frac{d}{2}-2}\Gamma\Big(\frac{d-2}{2}\Big)e^{\mathrm{i}\pi\frac{d-2}{2}}(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}\sum_{L=0}^{\infty}(2L+d-2)\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\mathsf{Q}^{\nu}_{k}(\sin\varphi_2)C^{\frac{d-2}{2}}_{L}(\cos\theta)\,.
\end{aligned} \qquad \text{(E.56)}$$

In particular, in even dimensions, we have

$$\begin{aligned}
&\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z)\\
&=2^{\frac{d}{2}-2}\,\Gamma\Big(\frac{d-2}{2}\Big)\,e^{\mathrm{i}\pi\frac{d-2}{2}}\,(\cos\varphi_1\,\cos\varphi_2)^{\frac{d-1}{2}}\,\sum_{L=0}^{\infty}(2L+d-2)\,\mathsf{P}^{-\nu}_{k}(\sin\varphi_1)\,\mathsf{Q}^{\nu}_{k}(\sin\varphi_2)\,C^{\frac{d-2}{2}}_{L}(\cos\theta)\,,
\end{aligned} \qquad \text{(E.57)}$$

which is equivalent to Eq. (E.50). Thus, when $Z<-1$, the right hand sides of Eqs. (E.25) and (E.26) always take the forms

$$\begin{cases}
0 & (d:\text{odd})\,,\\
\dfrac{\mathrm{i}}{(2\pi)^{\frac{d}{2}}\,\cos(\pi\nu)}\,e^{-\mathrm{i}\pi\frac{d-2}{2}}\,\big(Z^2-1\big)^{-\frac{d-2}{4}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-\frac{1}{2}}(-Z) & (d:\text{even})\,.
\end{cases} \qquad \text{(E.58)}$$

We thus have obtained simplified expressions for Eq. (E.30) for three different regions of Z in the form Eqs. (E.40), (E.46), and (E.58). One can readily see that three equations can be obtained from Eqs. (E.25) and (E.26). This completes the proof of our assertion.

F. Integral representation of the associated Legendre functions

In this appendix, we give a proof of Eq. (6.21) [we write it again here for convenience],

$$\mathcal{Q}^{\frac{d-2}{2}}_{\nu-1/2}(u) = e^{\mathrm{i}\pi\frac{d-2}{2}} \int_0^\infty \mathrm{d}\lambda\, \frac{\Gamma\big(\frac{d-1}{2}+\mathrm{i}\lambda\big)\,\Gamma\big(\frac{d-1}{2}-\mathrm{i}\lambda\big)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)}\, \frac{\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u)}{\nu^2+\lambda^2} \quad \big[d\in\mathbb{Z}\,, \quad \mathrm{Re}\,\nu>0\big]\,, \tag{F.1}$$

treating the odd and even dimensional cases separately. Our discussion is heavily based on the derivation of the heat kernel in Euclidean AdS space performed in [16] (see also [15]).

odd dimensions

When d is odd, using the direct relation between the associated Legendre functions and the Gegenbauer functions,

$$\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-\frac{1}{2}}(u) = \frac{2^{\frac{d-2}{2}}\,\pi^{1/2}\,\Gamma\big(\frac{d-1}{2}\big)}{\sin\big[\pi\,\big(\frac{d-1}{2}-\mathrm{i}\lambda\big)\big]\,\Gamma\big(\frac{d-1}{2}+\mathrm{i}\lambda\big)\,\Gamma\big(\frac{d-1}{2}-\mathrm{i}\lambda\big)}\,(u^2-1)^{\frac{d-2}{4}}\,C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)\,, \tag{F.2}$$

we can rewrite the integral on the right-hand side of (F.1) as

$$\begin{aligned}&\int_0^\infty \mathrm{d}\lambda\, \frac{\Gamma\big(\frac{d-1}{2}+\mathrm{i}\lambda\big)\,\Gamma\big(\frac{d-1}{2}-\mathrm{i}\lambda\big)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)}\, \frac{\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u)}{\nu^2+\lambda^2}\\ &= \frac{2^{\frac{d-2}{2}}\,e^{\mathrm{i}\pi d/2}\,\Gamma\big(\frac{d-1}{2}\big)}{\pi^{1/2}}\,\big(u^2-1\big)^{\frac{d-2}{4}} \int_0^\infty \mathrm{d}\lambda\,\frac{\lambda}{\nu^2+\lambda^2}\,C^{\frac{d-1}{2}}_{\mathrm{i}\lambda-\frac{d-1}{2}}(u)\,.\end{aligned} \tag{F.3}$$

By further using the identities for the Gegenbauer function

$$\mathrm{i}\lambda\,C^n_{\mathrm{i}\lambda-n}(\cosh\gamma) = \frac{2^{1-n}}{\Gamma(n)}\left[\frac{\mathrm{d}}{\mathrm{d}(\cosh\gamma)}\right]^n\cos(\lambda\gamma) \tag{F.4}$$

with nonnegative integers n, Eq. (F.3) can be rewritten as follows:

$$\begin{aligned}&= \frac{2^{1/2}\,e^{\mathrm{i}\pi\frac{d-1}{2}}}{\pi^{1/2}}\,\big(\cosh^2\gamma-1\big)^{\frac{d-2}{4}}\left[\frac{\mathrm{d}}{\mathrm{d}(\cosh\gamma)}\right]^{\frac{d-1}{2}}\int_0^\infty \mathrm{d}\lambda\,\frac{\cos(\lambda\gamma)}{\nu^2+\lambda^2}\\ &= \frac{\pi^{1/2}\,e^{\mathrm{i}\pi\frac{d-1}{2}}}{2^{1/2}\,\nu}\,\big(\cosh^2\gamma-1\big)^{\frac{d-2}{4}}\left[\frac{\mathrm{d}}{\mathrm{d}(\cosh\gamma)}\right]^{\frac{d-1}{2}} e^{\mp\gamma\nu} \quad [\,\mathrm{Re}\,\gamma\gtrless 0\,,\quad \mathrm{Re}\,\nu>0\,]\\ &= e^{-\mathrm{i}\pi\frac{d-1}{2}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-1/2}(\cosh\gamma) \qquad [\,\mathrm{Re}\,\nu>0\,]\,.\end{aligned} \tag{F.5}$$

Here, in order to show the second equality, we have used the formula [3.723-2] of [22]:

$$\int_0^\infty \mathrm{d}\lambda\,\frac{\cos(\lambda\gamma)}{\nu^2+\lambda^2} = \begin{cases} \dfrac{\pi}{2\nu}\,e^{-\gamma\,\nu} & [\,\mathrm{Re}\,\gamma>0\,,\quad \mathrm{Re}\,\nu>0\,] \\ \dfrac{\pi}{2\nu}\,e^{\gamma\,\nu} & [\,\mathrm{Re}\,\gamma<0\,,\quad \mathrm{Re}\,\nu>0\,] \end{cases}\,, \tag{F.6}$$

and to show the third equality we have used the following formula [see (A.20) of [16]]:

$$e^{-\mathrm{i}\pi\frac{d-2}{2}}\,\mathcal{Q}^{\frac{d-2}{2}}_{\nu-1/2}(\cosh\gamma) = \begin{cases} \dfrac{\pi^{1/2}\,e^{\mathrm{i}\pi\frac{d-1}{2}}}{2^{1/2}\,\nu}\,\bigl(\cosh^2\gamma-1\bigr)^{\frac{d-2}{4}}\left[\dfrac{\mathrm{d}}{\mathrm{d}(\cosh\gamma)}\right]^{\frac{d-1}{2}}e^{-\gamma\,\nu} & [\mathrm{Re}\,\gamma>0] \\ \dfrac{\pi^{1/2}\,e^{\mathrm{i}\pi\frac{d-1}{2}}}{2^{1/2}\,\nu}\,\bigl(\cosh^2\gamma-1\bigr)^{\frac{d-2}{4}}\left[\dfrac{\mathrm{d}}{\mathrm{d}(\cosh\gamma)}\right]^{\frac{d-1}{2}}e^{\gamma\,\nu} & [\mathrm{Re}\,\gamma<0] \end{cases}\,. \tag{F.7}$$

even dimensions

When d is even, the equality

$$\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-\frac{1}{2}}(u) = \frac{\Gamma\bigl(\mathrm{i}\lambda-\frac{d-1}{2}+1\bigr)}{\Gamma\bigl(\mathrm{i}\lambda+\frac{d-1}{2}\bigr)}\,\mathcal{P}^{\frac{d-2}{2}}_{\mathrm{i}\lambda-\frac{1}{2}}(u) \tag{F.8}$$

holds, and we can show (F.1) as follows:

$$\begin{aligned} &\int_0^\infty \mathrm{d}\lambda\,\frac{\Gamma\bigl(\frac{d-1}{2}+\mathrm{i}\lambda\bigr)\,\Gamma\bigl(\frac{d-1}{2}-\mathrm{i}\lambda\bigr)}{\Gamma(\mathrm{i}\lambda)\,\Gamma(-\mathrm{i}\lambda)}\,\frac{\mathcal{P}^{-\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u)}{\nu^2+\lambda^2} \\ &= (-1)^{\frac{d-2}{2}}\int_0^\infty \mathrm{d}\lambda\,\frac{\lambda\,\tanh(\pi\lambda)}{\nu^2+\lambda^2}\,\mathcal{P}^{\frac{d-2}{2}}_{\mathrm{i}\lambda-1/2}(u) \\ &= (-1)^{\frac{d-2}{2}}\bigl(u^2-1\bigr)^{\frac{d-2}{4}}\left(\frac{\mathrm{d}}{\mathrm{d}u}\right)^{\frac{d-2}{2}}\int_0^\infty \mathrm{d}\lambda\,\frac{\lambda\,\tanh(\pi\lambda)}{\nu^2+\lambda^2}\,\mathcal{P}_{\mathrm{i}\lambda-1/2}(u) \\ &= (-1)^{\frac{d-2}{2}}\bigl(u^2-1\bigr)^{\frac{d-2}{4}}\left(\frac{\mathrm{d}}{\mathrm{d}u}\right)^{\frac{d-2}{2}} Q_{\nu-1/2}(u) \\ &= e^{-\mathrm{i}\pi\frac{d-2}{2}}\,Q^{\frac{d-2}{2}}_{\nu-1/2}(u)\,, \end{aligned} \tag{F.9}$$

where we have used the identities for the associated Legendre functions with integer order,

$$\bigl(u^2-1\bigr)^{n/2}\,\frac{\mathrm{d}^n}{\mathrm{d}u^n}\,\mathcal{P}_{\mathrm{i}\lambda-\frac{1}{2}}(u) = \mathcal{P}^n_{\mathrm{i}\lambda-\frac{1}{2}}(u)\,,\quad \bigl(u^2-1\bigr)^{n/2}\,\frac{\mathrm{d}^n}{\mathrm{d}u^n}\,\mathcal{Q}_{\nu-\frac{1}{2}}(u) = \mathcal{Q}^n_{\nu-\frac{1}{2}}(u)\,, \tag{F.10}$$

and the formula [7.213] of [22]:

$$\int_0^\infty \mathrm{d}\lambda\,\frac{\lambda\,\tanh(\pi\lambda)}{\nu^2+\lambda^2}\,\mathcal{P}_{\mathrm{i}\lambda-1/2}(u) = Q_{\nu-1/2}(u) \qquad [\mathrm{Re}\,\nu>0]\,. \tag{F.11}$$

References

[1] T. S. Bunch and P. C. W. Davies, "Quantum Field Theory in de Sitter Space: Renormalization by Point Splitting," Proc. Roy. Soc. Lond. A **360**, 117 (1978).

[2] E. Mottola, "Particle Creation in de Sitter Space," Phys. Rev. D **31**, 754 (1985).

[3] B. Allen, "Vacuum States in de Sitter Space," Phys. Rev. D **32**, 3136 (1985).

[4] G. W. Gibbons and S. W. Hawking, "Cosmological Event Horizons, Thermodynamics, and Particle Creation," Phys. Rev. D **15**, 2738 (1977).

[5] R. Bousso, A. Maloney and A. Strominger, "Conformal vacua and entropy in de Sitter space," Phys. Rev. D **65**, 104039 (2002) [hep-th/0112218].

[6] A. AGrib, S. G. Mamaev and V. M. Mostepanenko, Gen. Rel. Grav. **7**, 535 (1976).

[7] N. D. Birrell and P. C. W. Davies, "Quantum Fields In Curved Space," Cambridge, UK: Univ. Pr. (1982).

[8] M. Castagnino and F. D. Mazzitelli, Phys. Rev. D **31**, 742 (1985).

[9] A. M. Polyakov, "De Sitter space and eternity," Nucl. Phys. B **797**, 199 (2008) [arXiv:0709.2899 [hep-th]].

[10] E. Alvarez and R. Vidal, "Eternity and the cosmological constant," JHEP **0910**, 045 (2009) [arXiv:0907.2375 [hep-th]].

[11] J. S. Schwinger, J. Math. Phys. **2**, 407 (1961).

[12] L. V. Keldysh, Zh. Eksp. Teor. Fiz. **47**, 1515 (1964) [Sov. Phys. JETP **20**, 1018 (1965)].

[13] D. Krotov and A. M. Polyakov, "Infrared Sensitivity of Unstable Vacua," Nucl. Phys. B **849**, 410 (2011) [arXiv:1012.2107 [hep-th]].

[14] A. M. Polyakov, "Gauge Fields And Strings," Contemp. Concepts Phys. **3**, CRC Press (1987).

[15] E. T. Akhmedov, "Real or Imaginary? (On pair creation in de Sitter space)," Mod. Phys. Lett. A **25**, 2815 (2010) [arXiv:0909.3722 [hep-th]].

[16] C. Grosche and F. Steiner, "The Path Integral On The Pseudosphere," Annals Phys. **182**, 120 (1988).

[17] I. R. Klebanov and E. Witten, "AdS / CFT correspondence and symmetry breaking," Nucl. Phys. B **556**, 89 (1999) [hep-th/9905104].

[18] A. Strominger, "The dS / CFT correspondence," JHEP **0110** (2001) 034 [hep-th/0106113].

[19] S. Weinberg, Phys. Rev. D **72**, 043514 (2005) [hep-th/0506236].

[20] M. Fukuma, Y. Sakatani and S. Sugishita, work in progress.

[21] M. Parikh, P. Samantray and E. Verlinde, "Rotating Rindler-AdS Space," Phys. Rev. D **86**, 024005 (2012) [arXiv:1112.3433 [hep-th]].

[22] I. S. Gradshteyn and I. M. Ryzhik, "Table of Integrals, Series, and Products," Alan Jeffrey and Daniel Zwillinger (eds.), Seventh edition (2007).

[23] A. Erdélyi et al., "Higher Transcendental Functions," Vol. I, McGraw-Hill Book Company, New York, (1953).

[24] N. N. Beloozerov, "Addition theorems for Legendre functions as corollaries of an addition theorem for Gegenbauer polynomials," Mathematical notes of the Academy of Sciences of the USSR, Volume 20, Issue 3, pp 740-745 (1976).